



H·CUP

HEALTHCARE COST AND UTILIZATION PROJECT

HCUP Methods Series



Agency for Healthcare
Research and Quality



U.S. Department of Health and Human Services
Agency for Healthcare Research and Quality

**Contact Information:
Healthcare Cost and Utilization Project (HCUP)
Agency for Healthcare Research and Quality
5600 Fishers Lane
Room 07W17B
Mail Stop 7W25B
Rockville, MD 20857
<http://www.hcup-us.ahrq.gov>**

For Technical Assistance with HCUP Products:

Email: hcup@ahrq.gov

or

Phone: 1-866-290-HCUP

Recommended Citation: Barrett M, Coffey R, Houchens R, Moy E, Heslin K, Moles E, Coenen N. *Methods Applying AHRQ Quality Indicators to Healthcare Cost and Utilization Project (HCUP) Data for the 2015 National Healthcare Quality and Disparities Report (QDR)*. 2016. HCUP Methods Series Report # 2016-01 ONLINE. April 12, 2016. U.S. Agency for Healthcare Research and Quality. Available: <http://www.hcup-us.ahrq.gov/reports/methods/methods.jsp>.

TABLE OF CONTENTS

INTRODUCTION	1
AHRQ QUALITY INDICATORS.....	1
PREPARATION OF HCUP DATABASES	2
HCUP Data on Race/Ethnicity.....	2
Modifications to the HCUP Databases	3
STEPS TAKEN TO APPLY AHRQ QUALITY INDICATORS TO THE HCUP DATA.....	4
Review and Modify the AHRQ QI Software	4
Acquire Population-Based Data for Denominators and Risk-Adjustment.....	4
Assign QI Indicators to the HCUP Databases	5
Adapt Statistical Methods to HCUP Data	5
SPECIAL ANALYSES	7
Trends in National Inpatient Hospital Costs Associated with Quality Indicators	7
Trends in IQI and PSI Summary Measures	7
Benchmarks for State Performance for the Quality Indicators	7
Trends in National Inpatient Stays with Multiple Chronic Conditions	7
Medicaid and Uninsured Inpatient Stays and Aggregate Hospital Costs in the United States.....	8
Inpatient Stays for Select Conditions Treated near the United States-Mexico Border.....	8
Focus on Select PQIs by Race/Ethnicity in Hawaii.....	8
National Estimates of Emergency Department Utilization.....	9
TABLES	10
Table 1. AHRQ Quality Indicators Applied to the HCUP Data for the National Healthcare Quality and Disparities Report (QDR).....	11
Table 2. Sources of 2013 HCUP Inpatient Data for the QDR	21
Table 3. HCUP National and State Inpatient Databases Used for QI Estimates	23
Table 4. Age Groupings for Risk Adjustment.....	27
REFERENCES	28
APPENDIX A: DEVELOPMENT OF NATIONALLY WEIGHTED ANALYSIS FILES FOR THE NATIONAL HEALTHCARE QUALITY AND DISPARITIES REPORT	1
2013 Nationally Weighted Analysis File for the QDR.....	1
Preparing the 2013 Nationally Weighted QDR Analysis File.....	2
Evaluating the 2013 Nationally Weighted QDR Analysis File	3
APPENDIX B: MODIFICATION TO THE STATE INPATIENT DATABASES FOR STATE- LEVEL REPORTING BY RACE/ETHNICITY	1
APPENDIX C: EMERGENCY DEPARTMENT RATES FOR SELECTED AHRQ QUALITY INDICATORS	1
APPENDIX D: STATISTICAL METHODS	1
APPENDIX E: CAVEATS TO THE INTREPRETAION OF HCUP-BASED QI ESTIMATES REPORTED IN THE NATIONAL HEALTHCARE QUALITY AND DISPARITIES REPORT	1

ICD-9-CM Coding Changes	1
Data Collection Differences Among States.....	1
Data Elements Used for QI Exclusions.....	1
Number of Clinical Fields	2
Use of External Cause-of-Injury Codes	4
Non-Resident Discharges in State-Level Estimates	4
Variation Among State QI Rates	5
APPENDIX F: QDR SUMMARY MEASURES FOR PATIENT SAFETY AND MORTALITY FOR SELECTED PROCEDURES AND CONDITIONS	1
IQIs and PSIs Used for the QDR Summary Measures	1
Calculation of the IQI and PSI Summary Measures.....	3

INTRODUCTION

The Agency for Healthcare Research and Quality (AHRQ) Quality Indicators (QIs) were applied to the Healthcare Cost and Utilization Project (HCUP) hospital discharge data for selected measures in the National Healthcare Quality and Disparities Report (QDR).¹ The report measures and tracks trends in quality and disparities in six key areas of health care that align with the National Quality Strategy (NQS) priorities — patient safety, person-centered care, care coordination, effective treatment, healthy living, and care affordability — and access to health care, which cuts across the priorities. A focus on priority populations summarizes quality and disparities in care for populations at elevated risk for receiving poor health care, which includes HCUP-based measures related to racial, ethnic, and socioeconomic factors for priority populations, including changes over time and across the urban-rural continuum. The QDR provides a comprehensive overview of the quality of health care received by the general population and disparities in care experienced by different racial, ethnic, and socioeconomic groups. Electronic dissemination of the report was expanded and information on individual measures will be available online through a series of chartbooks.²

This report describes the preparation of the HCUP databases for data years 2000–2013 for use in the 2015 QDR, the steps taken to apply the AHRQ QIs to the HCUP data, and other analyses based on HCUP data that are not specific to the QIs, but are developed for use in the 2015 QDR.

AHRQ QUALITY INDICATORS

The AHRQ QIs are measures of quality associated with processes of care that occur in an outpatient or an inpatient setting. The QIs rely solely on hospital inpatient administrative data and, for this reason, are screens for examining quality that may indicate the need for more in-depth studies. The AHRQ QIs used for the QDR include four sets of measures:

- Prevention Quality Indicators (PQIs) — or ambulatory care sensitive conditions — identify hospital admissions that evidence suggests could have been avoided, at least in part, through high-quality outpatient care (AHRQ, 2012).
- Inpatient Quality Indicators (IQIs) reflect quality of care inside hospitals and include measures of utilization of procedures for which there are questions of overuse, underuse, or misuse (AHRQ, 2012).
- Patient Safety Indicators (PSIs) reflect quality of care inside hospitals, by focusing on surgical complications and other iatrogenic events (AHRQ, 2012).
- Pediatric Quality Indicators (PDIs) reflect quality of care inside hospitals and identify potentially avoidable hospitalizations among children (AHRQ, 2012).

The AHRQ QI measures generated for possible inclusion in the QDR are described in [Table 1](#) at the end of this methods report. Not all of these QIs were used in the reports. The 2015 QDR includes QI trends through data year 2013. The following national and State-level QI estimates were constructed from the HCUP databases for the QDR:

¹Beginning with the 2014 report, findings that previously appeared in two separate reports (the *National Healthcare Quality Report* and the *National Healthcare Disparities Report*) have been integrated into a single document — the National Healthcare Quality and Disparities Report (QDR).

² The National Healthcare Quality & Disparities Reports and Chartbooks are available at <http://www.ahrq.gov/research/findings/nhqdr/index.html>.

- National inpatient trends by QI, overall and by subpopulations including community income, expected primary payer, and race/ethnicity
- State-level inpatient trends by QI, overall and by subpopulations including community income, expected primary payer, and race/ethnicity, for HCUP Partner organizations that agreed to participate in the QDR
- National trends for selected PQIs and PDIs in the emergency department setting.

PREPARATION OF HCUP DATABASES

The Healthcare Cost and Utilization Project (HCUP) is a family of healthcare databases and related software tools and products developed through a Federal-State-Industry partnership and sponsored by AHRQ. HCUP databases are derived from administrative data and contain encounter-level, clinical and nonclinical information including all-listed diagnoses and procedures, discharge status, patient demographics, and charges for all patients, regardless of payer (e.g., Medicare, Medicaid, private insurance, uninsured), beginning in 1988. These databases enable research on a broad range of health policy issues, including cost and quality of health services, medical practice patterns, patient safety, access to health care programs, and outcomes of treatments at the national, State and local market levels.

The following HCUP databases were used as the source of data for the QDR:

- The HCUP Nationwide Inpatient Sample (NIS), a nationally stratified sample of hospitals (with all of their discharges) from States that contribute data to HCUP for trends in data years 2000–2011. A similarly constructed nationally stratified sample of hospitals from HCUP States for trends in data years 2012–2013³
- The HCUP State Inpatient Databases (SID), a census of hospitals (with all of their discharges) from participating States in data years 2004 and 2011–2013
- The HCUP Nationwide Emergency Department Sample (NEDS), a nationally stratified sample of hospital-based emergency departments (with information for both treat-and-release visits and those resulting in a hospital admission) data years 2008–2013.

The list of HCUP Partner organizations that contribute to the HCUP databases is provided in [Table 2](#) at the end of this methods report. The HCUP databases used for each year’s national inpatient estimates are described in [Table 3](#).

HCUP Data on Race/Ethnicity

HCUP coding includes race and ethnicity in one data element (RACE). Because of variability in the collection of race and ethnicity information in the State data, HCUP maintains a uniform set of categories based on race definitions used in the 1977 Office of Management and Budget (OMB) Directive 15 using the combined race-ethnicity format (separate categories for Hispanic and five Non-Hispanic racial groups – White, Black, Asian or Pacific Islander, American Indian or Alaska Native (AIAN), and Other).

³ In data year 2012, the HCUP NIS changed its design and became a nationally stratified sample of discharges. The definition of the discharge universe was also revised to exclude long-term acute care hospitals. Because of the differences in design, the trend analyses for the QDR did not use the 2012–2013 NIS.

When a State and its hospitals collect Hispanic ethnicity *separately* from race, HCUP assigns the data to the combined race/ethnicity categorization and uses Hispanic ethnicity to override any other race category to create uniform coding across States.

There is also limited reporting of AIAN in the HCUP data. In addition, in some areas of the country care for the AIAN population is provided in Indian Health Service (IHS) hospitals, which are not included in HCUP. For these reasons, AIAN discharges were combined with “Other” races for the QDR analyses.

The resulting QDR reporting categories for the HCUP data include: White Non-Hispanic; African American Non-Hispanic; Asian/Pacific Islander Non-Hispanic; Other Non-Hispanic; and Hispanic (of any race).

Modifications to the HCUP Databases

In preparation for the QDR and its derivative products, the HCUP databases needed to be customized as indicated below:

1. The HCUP SID were modified to create analytic files consistent across States.
 - *Subset to Community Hospitals.* For the SID, we selected community hospitals⁴ and eliminated rehabilitation hospitals.
 - *Weight for Missing Hospitals.* Because some statewide data organizations do not report data for all community hospitals in the State, we weighted hospitals in the SID to the State’s universe of hospitals in the American Hospital Association (AHA) Annual Survey Database based on hospital characteristics.
 - *Weight for Missing Quarters.* Discharges from hospitals operating for the entire year but not contributing data for one or more quarters were weighted up to annual estimates for that institution in the SID.
2. The HCUP databases were augmented as necessary for the QDR analyses:
 - *Impute for Missing Characteristics.* For missing age, sex, race/ethnicity, ZIP Code, and expected primary payer data that occurred on a small proportion of discharge records, we used a “hot deck” imputation method (which draws donors from strata of similar hospitals and patients) to assign values while preserving the variance within the data.
 - *Assign Additional Measures for Reporting.* We assigned median household income quartile by linking Nielsen ZIP Code demographic data to patient’s ZIP Code in the SID. Income quartiles were defined annually based on the distribution of the population in the United States.

⁴ *Community* hospitals are defined by the AHA as “non-Federal, short-term, general, and other specialty hospitals, excluding hospital units of institutions.” The specialty hospitals included in the AHA definition of “community hospitals” are: obstetrics-gynecology, ear-nose-throat, short-term rehabilitation, orthopedic, and pediatric institutions. The AHA also groups public hospitals and academic medical centers with community hospitals. Starting in 2005, the AHA included long term acute care facilities in the definition of community hospitals, therefore such facilities are included in the NIS sampling frame. These facilities provide acute care services to patients who need long term hospitalization (stays of more than 25 days). Excluded from the AHA definition of “community hospitals” are long-term non-acute care hospitals, psychiatric hospitals, and alcoholism/chemical dependency treatment facilities. For the QDR analyses, we selected all AHA-defined “community hospitals” with the exception of short-term rehabilitation hospitals (beginning with 1998 HCUP data).

3. The HCUP SID were used to create QDR analysis files designed to provide national estimates for overall and priority population reporting (for all groups including race/ethnicity). [Appendix A](#) to this report provides detail on the creation of the QDR analysis files for national estimates. The SID included in each of the QDR analysis files are listed in [Table 3](#).
4. The HCUP SID were also used for reporting overall and by priority populations within State (including community income quartile, expected primary payer, and race/ethnicity). State-level QI estimates are only reported for participating HCUP Partners that agree to release information in the QDR. Reporting by race/ethnicity was limited to SID that included this type of information. [Appendix B](#) to this report provides additional detail on the preparation of the SID for reporting by race/ethnicity.
5. The NEDS were used to calculate PQIs and PDIs in the emergency department setting. A description of the data preparation and methods used for national QI estimates from the NEDS is included in [Appendix C](#).

STEPS TAKEN TO APPLY AHRQ QUALITY INDICATORS TO THE HCUP DATA

To apply the AHRQ Quality Indicators to HCUP hospital discharge data for the QDR, several steps were taken: (1) QI software review and modification, (2) acquisition of population-based data, (3) assignment of QIs to the HCUP databases, and (4) identification of statistical methods.

Review and Modify the AHRQ QI Software. For the 2015 QDR, we started with the following QI software versions: PQI Version 4.4, IQI Version 4.4, PSI Version 4.4, and PDI Version 4.4. Because each of these software modules was developed for State and hospital-level rates, rather than national rates, the additional step of weighting the QI estimates was necessary.

In addition, we did not utilize the present on admission (POA) estimation module for the IQIs, PDIs, and PSIs since POA indicators were not uniformly available from States that contribute to the HCUP databases. Other QI-specific modifications are noted as footnotes in the tables.

We added three indicators particularly relevant to the structure of the QDR. One indicator was created for discharges age 65 years and older: immunization-preventable influenza, age 65 and over. Two additional indicators were created to facilitate longitudinal analyses by modifying the chronic and overall PQI composite measures to exclude PQI 05 for chronic obstructive pulmonary disease (COPD) and asthma for patients aged 40 years and older. Because of ICD-9-CM coding changes for COPD for data prior to 2005 rates are not compatible with rates for 2005 forward.

Acquire Population-Based Data for Denominators and Risk-Adjustment. The next step was to acquire data for the numerator and denominator populations for the QIs. The AHRQ QIs measure an event that occurs in a hospital, requiring a numerator count of the event of interest and a denominator count of the population (within a hospital or geographic area) to which the event relates.

For the numerator counts of the AHRQ QIs, we used the HCUP databases. For the denominator counts, we identified two sources for all reporting categories and for all adjustment categories listed in the HCUP-based tables.

- For QIs that related to *providers*, the HCUP data were used for national and State-level discharge denominator counts.

- For QIs that related to *geographic areas*, population ZIP-Code-level counts from demographic update data provided by Nielsen (a vendor that compiles and adds value to the U.S. Bureau of Census data) were used for denominator counts. Nielsen uses intra-census methods to estimate household and demographic statistics for geographic areas (The Nielsen Company). We also used the Nielsen population data for risk adjustment by age and sex for the area-based QIs.

Assign QI Indicators to the HCUP Databases. The four AHRQ QI program modules were applied to the prepared SID data using all available diagnoses and procedures reported by each State. The QI indicators from the SID were then linked to the corresponding discharge records on the 2000–2011 NIS.

Adapt Statistical Methods to HCUP Data. Several statistical issues needed to be addressed when applying the AHRQ QI software to the HCUP data, including: age-sex adjustment for all QIs; severity/comorbidity adjustment for the discharge-based IQIs, PSIs, and PDIs; and derivation of standard errors and appropriate hypothesis tests.

- *Age-Sex Risk Adjustment.* For the *PQIs* and area-based *IQIs*, *PSIs*, and *PDIs*, the observed rates were risk-adjusted for age and sex differences across population subgroups and were based on methods of direct standardization (Fleiss, 1973). Age was categorized into 18 five-year increments (described in [Table 4](#), Age Groupings for Risk Adjustment). Although the AHRQ QI software uses a similar approach to adjust the area-based QIs, we relied on direct standardization because of the additional reporting categories and denominators for priority populations required in the QDR.
- *Age, Sex, Severity, and Comorbidity Risk Adjustment.* For the discharge-based *PSIs*, the observed rates were risk-adjusted for age, sex, age-sex interaction, diagnosis related groups (DRG) cluster, and comorbidities using the regression-based standardization that is part of the AHRQ PSI software, with the following exceptions:
 - When reporting by age, the risk adjustment includes all of the above except age.
 - When reporting by sex, the risk adjustment includes all of the above except sex.

For the discharge-based *IQIs*, risk adjustments were made for age, sex, age-sex interaction, and the 3M™ All Patient Refined Diagnosis Related Groups (APR-DRGs) risk of mortality or severity score using the regression-based standardization that is part of the AHRQ IQI software, with the following exceptions:

- When reporting by age, the risk adjustment includes all of the above except age.
- When reporting by sex, the risk adjustment includes all of the above except sex.

For the discharge-based *PDIs*, risk adjustments were made for age, sex, DRG and major diagnostic category (MDC) clusters, and comorbidities using the regression-based standardization that is part of the AHRQ PDI software. Measure-specific stratification by risk group, clinical category, and procedure type was also applied, with the following exceptions:

- When reporting by age, the risk adjustment includes all of the above except age.
- When reporting by sex, the risk adjustment includes all of the above except sex.

- *Standard Errors and Hypothesis Tests.* Standard error calculations for the rates were based on the HCUP report entitled *Calculating Nationwide Inpatient Sample (NIS) Variances* (Houchens, et al., 2005). There is no sampling error associated with Nielsen

census population counts. Appropriate statistics for a stratified sample were obtained through the Statistical Analysis System (SAS) procedure called PROC SURVEYMEANS.

- *Masking Rates for Statistical Reliability, Data Quality, and Confidentiality.* QI estimates were included in the QDR if they reached a threshold defined by a relative standard error less than 30 percent and at least 11 unweighted cases in the denominator. Estimates that did not satisfy these criteria were masked (set to DSU, for “data statistically unreliable”). Statistical calculations are explained in [Appendix D](#) to this report.

Some caution should be used in interpreting the HCUP-based AHRQ QI statistics presented in the QDR. Limitations that relate to how the QIs were applied, ICD-9-CM coding changes over time, inter-State differences in data collection, and other issues are explained in detail in [Appendix E](#) to this report.

SPECIAL ANALYSES

Trends in National Inpatient Hospital Costs Associated with Quality Indicators

The QDR includes trends in total national costs for the select PQIs and PDIs including the composites and ambulatory care sensitive conditions such as heart failure, diabetes, and asthma. Total national costs associated with these PQIs and PDIs were calculated overall and by community income quartile and race/ethnicity.

Total charges were converted to costs using the hospital-level HCUP Cost-to-Charge Ratio Files (CCR Files) based on Hospital Cost Report data from the Centers for Medicare & Medicaid Services (CMS).⁵ Costs reflect the actual costs of production, while charges represent what the hospital billed for the stay. Hospital charges reflect the amount the hospital charged for the entire hospital stay and do not include professional (physician) fees. The total cost is the product of the number of stays for each QI measure and the mean cost for each QI measure. This approach compensates for stays for which charges (and thus estimated costs) are not available. Costs were adjusted to 2013 dollars for all years using the price indices for the gross domestic product (downloaded from the Bureau of Economic Analysis, U.S. Department of Commerce).

Trends in IQI and PSI Summary Measures

To examine national and State-level trends in inpatient mortality and patient safety events, risk-adjusted rates for select IQIs and PSIs were summarized. The three QDR summary measures include: (1) Mortality for *selected conditions* based on select IQIs; (2) Mortality for *selected procedures* based on select IQIs; and (3) Patient Safety based on select PSIs. These summary measures were calculated as a weighted sum of risk-adjusted rates for individual IQIs and PSIs. Additional information on the calculation of IQI and PSI Summary Measures is provided in [Appendix F](#).

Benchmarks for State Performance for the Quality Indicators

Based on a recommendation from the Institute of Medicine's report on *Future Directions for the National Healthcare Quality and Disparities Reports*, benchmarks based on a straight average of the top 10 percent of reporting States were determined. For a benchmark to be calculated, rates for at least 30 States needed to be available.

Trends in National Inpatient Stays with Multiple Chronic Conditions

The criteria for multiple chronic conditions (MCC) were developed by the Department of Health and Human Services (HHS) Interagency Workgroup on MCC and the Office of the Assistant Secretary of Health.⁶ The 20 different chronic conditions were identified using the AHRQ Clinical Classification Software (CCS)⁷ unless noted: hypertension (CCS 98–99), hyperlipidemia (CCS 53), congestive heart failure (CCS 108), coronary artery disease (CCS 100–101),

⁵ HCUP Cost-to-Charge Ratio Files. Healthcare Cost and Utilization Project (HCUP). September 2015. Agency for Healthcare Research and Quality, Rockville, MD. Available: www.hcup-us.ahrq.gov/db/state/costtocharge.jsp.

⁶ Goodman RA, Posner SF, Huang ES, Parekh AK, Koh HK. Defining and measuring chronic conditions: imperatives for research, policy, program and practice. *Prev Chronic Dis* 2013;10: 120239.

⁷ Steiner CA, Friedman B. Hospital Utilization, Costs, and Mortality for Adults With Multiple Chronic Conditions, Nationwide Inpatient Sample, 2009. *Prev Chronic Dis* 2013;10:120292.

diabetes (CCS 49–50), stroke (CCS 109–112), cardiac arrhythmias (CCS 105–106), arthritis (CCS 202–203), cancer (CCS 11–43), depression (CCS 657), dementia includes Alzheimer’s and other senile dementias (CCS 653), substance abuse disorders (CCS 660–661), chronic obstructive pulmonary disease (CSS 127), asthma (CCS 128), chronic kidney disease (CCS 156–158), HIV (CCS 5), hepatitis (CCS 6), autism spectrum disorder (ICD-9-CM diagnoses 29900, 29901), schizophrenia (CCS 659), and osteoporosis (CCS 206). Chronic conditions may be reported as a principal or secondary diagnosis. A specific chronic condition is only counted once per discharge. For example, if a discharge has one secondary diagnosis for hypertension included in CCS 98 and a different secondary diagnosis indicating hypertension included in CCS 99, the chronic condition of hypertension is only counted once for that discharge.

National trends from 2002–2013 in adult inpatient stays by the number of MCCs were developed using the HCUP National (Nationwide) Inpatient Sample (NIS). The NIS trend weights were used for data years 2002–2011 for consistent estimates over the entire study period.⁸

Medicaid and Uninsured Inpatient Stays and Aggregate Hospital Costs in the United States

Information on Medicaid and uninsured inpatient stays and aggregate hospital costs in the United States for 2012–2013 were developed using the HCUP NIS. Medicaid and uninsured discharges were identified based on the expected primary payer of Medicaid, self-pay, no charge, and charity. Discharge counts and aggregate hospital costs were reported as a percentage of the total U.S. and by hospital characteristics including region, ownership, teaching status, urban-rural location, and size of the hospital based on the number of beds.

Inpatient Stays for Select Conditions Treated near the United States-Mexico Border

The QDR, in collaboration with the US-Mexico Border Health Commission,⁹ compared inpatient stays related to priority conditions in counties along the U.S.-Mexico border to non-border U.S. counties in four States (Arizona, California, New Mexico, and Texas). Border counties were identified as being within 60 miles of Mexico. Discharges were grouped by the county of the hospital. The analysis was limited to community hospitals excluding rehabilitation hospitals and included all patients treated in those hospitals, including foreign and out-of-State patients. Results were reported by Hispanic ethnicity. Information was included in the QDR [Chartbook for Hispanic Health Care](#).

Focus on Select PQIs by Race/Ethnicity in Hawaii

The HCUP SID for Hawaii includes detailed reporting of the patient’s race/ethnicity, including the identification of Hawaiian, Chinese, Filipino, Japanese, Korean, and Samoan which are often included in the general category of Asian/Pacific Islander. Corresponding population data were obtained from the annual Hawaii Health Survey (HHS) collected by the Office of Health Status Monitoring, Hawaii Department of Health. PQI estimates for asthma, congestive heart failure, diabetes, pneumonia, and urinary tract information were developed for 2011–2012 using the more detailed race/ethnicity categories.

⁸ More information on the NIS trend weights is available on the HCUP User Support Web site at <http://www.hcup-us.ahrq.gov/db/nation/nis/trendwghts.jsp>.

⁹ More information on the US-Mexico Border Health Commission is available at http://www.borderhealth.org/healthy_border.php?curr=bhc_initiatives.

National Estimates of Emergency Department Utilization

Beginning in the 2009 NHQR, the HCUP Nationwide Emergency Department Sample (NEDS) was used to examine national and regional differences in emergency department (ED) use for selected PQIs and PDIs associated with ambulatory care sensitive conditions. Age-sex adjusted rates are calculated from 2008 forward for all ED visits and those that do and do not result in an inpatient admission. Details on the use of the NEDS for reporting by QI are provided in [Appendix C](#).

There were three additional condition-specific analyses based on the NEDS:

- Starting with data year 2007, the NEDS was used to examine national trends in ED visits for mental illness and substance use disorders. ED visits were identified by the CCS category for the first-listed diagnosis. CCS 650–659, 662, and 670 were used to define mental illness disorders. CCS 660–661 defined substance use disorders. No distinction was made between ED visits that resulted in a hospital admission and those that did not. Nielsen population data was used to calculate rates per 100,000 residents by age, sex, community income, urban-rural location of patient residence, and region of the United States. Rates were not risk-adjusted.
- Starting with data year 2010, the NEDS was used to examine national trends in ED visits for dental conditions. ED visits were identified by an ICD-9-CM principal diagnosis of 520.00–523.9. No distinction was made between ED visits that resulted in a hospital admission and those that did not. Nielsen population data was used to calculate rates per 100,000 residents by age, sex, community income, and urban-rural location of patient residence. Rates were not risk-adjusted.
- Starting with data year 2011, the NEDS was used to examine national trends in ED visits for severe injuries. Severity of injuries was determined using the ICD Programs for Injury Categorization (ICDPIC).¹⁰ The ICDPIC translates ICD-9-CM diagnosis codes into several standard injury categories and/or scores, including Injury Severity Scores (ISS). The ISS are an anatomical scoring system that provides an overall score for patients with multiple injuries. Each injury is assigned Abbreviated Injury Scale (AIS) scores that are allocated to six body regions: Head, Face, Chest, Abdomen, Extremities (including pelvis), and External. AIS scores are: 1 (minor), 2 (moderate), 3 (serious), 4 (severe), 5 (critical), and 6 (not survivable). The highest AIS score in each body region is recorded. Scores from the three most severely injured body regions are squared and summed to produce the ISS. The ISS correlates linearly with mortality, morbidity, hospital stay, and other measures of severity. Some severely-injured patients are stabilized in the ED; others require hospitalization for further care. A high ISS score does not necessarily indicate the need for admission. Injuries with an ISS of 16 or greater were considered severe. An ISS of 16 or greater was assigned when at least one injury had an AIS score of 4 (severe) or at least two injuries had an AIS score of 3 (serious). ED utilization for severe injuries was reported by trauma level (trauma level I and II combined, trauma level III, and non-trauma) and by age, sex, community income, urban-rural location of patient residence, and region of the United States.

¹⁰ The ICDPIC does not calculate a severity score for late effects of injuries (ICD-9-CM diagnoses 905–909), effects of foreign body (930-939), burns (940–949), certain early complications of trauma (958), or poisoning by drugs, toxins, and other effects (960–995).

TABLES

Table 1. AHRQ Quality Indicators Applied to the HCUP Data for the National Healthcare Quality and Disparities Report (QDR)

This table includes the list of all version 4.4 AHRQ Quality Indicators (QIs) calculated using HCUP data. Not all of the AHRQ QIs listed below were included in the 2015 QDR.

QI No.	Description	Footnote
Prevention Quality Indicators¹¹		
PQI 1	Admissions with diabetes with short-term complications ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PQI software, diabetes must be the principal diagnosis and short-term complications include ketoacidosis, hyperosmolality, or coma. Transfers from other institutions are excluded.
PQI 2	Admissions with perforations or abscesses of appendix per 1,000 admissions with appendicitis ^a , age 18 and over	^a Consistent with the AHRQ PQI software, obstetric discharges and transfers from other institutions are excluded.
PQI 3	Admissions with diabetes with long-term complications ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PQI software, diabetes must be the principal diagnosis and long-term complications include renal, eye, neurological, circulatory, or other unspecified complications. Transfers from other institutions are excluded.
PQI 5	Admissions with chronic obstructive pulmonary disease (COPD) ^a or asthma per 100,000 population, age 40 and over	^a Consistent with the AHRQ PQI software, the principal diagnosis must be COPD, asthma, or acute bronchitis with COPD as a secondary diagnosis. Transfers from other institutions are excluded.
PQI 7	Admissions with hypertension ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PQI software, hypertension must be the principal diagnosis and exclusions include the following: admissions with kidney disease with dialysis access procedures, admissions with cardiac procedures, and transfers from other institutions.
PQI 8	Admissions for heart failure (HF) ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PQI software, HF must be the principal diagnosis and exclusions include the following: admissions with cardiac procedures and transfers from other institutions.
PQI 9	Low birth weight infants per 1,000 newborns ^a	^a Consistent with the AHRQ PQI software, exclusions include transfers from other institutions.
PQI 10	Admissions for dehydration ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PQI software, dehydration may be a principal diagnosis or a secondary diagnosis with a principal diagnosis of hyperosmolality and/or hypernatremia, gastroenteritis, or acute kidney injury. Exclusions include the following: admissions with a diagnosis code for chronic renal failure and transfers from other institutions.
PQI 11	Admissions for bacterial pneumonia ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PQI software, bacterial pneumonia must be the principal diagnosis and exclusions include the following: admissions for sickle cell disease or HB-S disease, admissions in an immunocompromised state, and transfers from other institutions.
PQI 12	Admissions for urinary tract infection (UTI) ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PQI software, UTI must be the principal diagnosis and exclusions include the following: admissions with kidney or urinary tract disorders, admissions in an immunocompromised state, and transfers from other institutions.
PQI 13	Admissions for angina without cardiac procedure ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PQI software, angina must be the principal diagnosis, and exclusions include admissions with cardiac procedures and transfers from other institutions.

¹¹ Indicators PQI 4 and PQI 6 are not assigned by the PQI software, version 4.4.

QI No.	Description	Footnote
PQI 14	Admissions for uncontrolled diabetes without complications ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PQI software, diabetes without complications must be the principal diagnosis and exclusions include transfers from other institutions.
PQI 15	Admissions for asthma ^a per 100,000 population, age 18 to 39	^a Consistent with the AHRQ PQI software, asthma must be the principal diagnosis on admissions ages 18 to 39 years old, and the following cases are excluded: admissions with cystic fibrosis or anomalies of the respiratory system and transfers from other institutions.
PQI 16	Lower extremity amputations among admissions for diabetes ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PQI software, a procedure code for lower-extremity amputation and a diagnosis of diabetes must be present. Exclusions include admissions for toe amputation or traumatic amputations of the lower extremity, obstetric discharges, and transfers from other institutions.
PQI 17 (Added)	Admissions for immunization-preventable pneumococcal pneumonia ^a per 100,000 population, age 65 and over	^a Immunization-preventable pneumococcal pneumonia may be reported as either the principal diagnosis or a secondary diagnosis. Exclusions include transfers from other institutions.
PQI 18 (Added)	Admissions for immunization-preventable influenza ^a per 100,000 population, age 65 and over	^a Immunization-preventable influenza may be reported as either the principal diagnosis or a secondary diagnosis. Exclusions include transfers from other institutions.
PQI 90	AHRQ overall Prevention Quality Indicator (PQI) composite ^a per 100,000 population, age 18 and over	^a Based on the twelve AHRQ PQIs for angina, asthma, bacterial pneumonia, chronic obstructive pulmonary disease, congestive heart failure, dehydration, diabetes, hypertension, and urinary tract infection.
PQI 90x (Added)	AHRQ modified ^a overall Prevention Quality Indicator (PQI) composite per 100,000 population, age 18 and over	^a Based on the eleven AHRQ PQIs for angina, asthma, bacterial pneumonia, congestive heart failure, dehydration, diabetes, hypertension, and urinary tract infection. For consistency of longitudinal reporting, the modified overall composite does not include AHRQ PQI 5 for chronic obstructive pulmonary disease because it is affected by ICD-9-CM coding changes.
PQI 91	AHRQ acute Prevention Quality Indicator (PQI) composite ^a per 100,000 population, age 18 and over	^a Based on the three AHRQ PQIs for bacterial pneumonia, dehydration, and urinary tract infection.
PQI 92	AHRQ chronic Prevention Quality Indicator (PQI) composite ^a per 100,000 population, age 18 and over	^a Based on the nine AHRQ PQIs for angina, asthma, chronic obstructive pulmonary disease, congestive heart failure, diabetes, and hypertension.
PQI 92x (Added)	AHRQ modified ^a chronic Prevention Quality Indicator (PQI) composite per 100,000 population, age 18 and over	^a Based on the eight AHRQ PQIs for angina, asthma, congestive heart failure, diabetes, and hypertension. For consistency of longitudinal reporting, the modified overall composite does not include AHRQ PQI 5 for chronic obstructive pulmonary disease because it is affected by ICD-9-CM coding changes.
Pediatric Quality Indicators¹²		
PDI 01	Admissions with accidental puncture or laceration during procedure per 1,000 medical and surgical admissions, ^a age less than 18 years	^a The AHRQ PDI software requires that the accidental puncture or laceration be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PDI software, the secondary diagnosis could be present on admission. Consistent with the AHRQ PDI software, the following cases are excluded: obstetric admissions, admissions involving spinal surgery, normal newborns, and neonates with a birth weight less than 500 grams.

¹² Indicator PDI 4 is not assigned by the PDI software, version 4.4. Incidence measures PDI 2 (pressure ulcer), PDI 3 (foreign body), and PDI 13 (transfusion reaction) are not calculated. Volume measure PDI 7 (pediatric heart surgery) is also not calculated.

QI No.	Description	Footnote
PDI 05	Admissions with iatrogenic pneumothorax per 1,000 medical and surgical admissions, ^a age less than 18 years	^a The AHRQ PDI software requires that the iatrogenic pneumothorax be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PDI software, the secondary diagnosis could be present on admission. Consistent with the AHRQ PDI software, the following cases are excluded: obstetric admissions, normal newborns, neonates with a birth weight less than 2500 grams, and admissions with chest trauma, pleural effusion, thoracic surgery, lung/pleural biopsy, diaphragmatic surgery repair, or cardiac surgery.
PDI 06	Deaths per 1,000 heart surgery admissions, ^a age less than 18 years	^a Consistent with the AHRQ PDI software, exclusions include obstetric admissions; admissions with transcatheter interventions as a single cardiac procedure or performed without bypass, but with catheterization; admissions with septal defects as single cardiac procedures without bypass; admissions with an atrial septal defect or ventricular septal defect repair with patent ductus arteriosus as the only cardiac procedure; heart transplants; premature infants with patent ductus arteriosus (PDA) closure as only cardiac procedure; infants age less than 30 days with PDA closure as only cardiac procedure; transfers to another hospital; and neonates with a birth weight less than 500 grams.
PDI 08	Postoperative hemorrhage or hematoma with surgical drainage or evacuation per 1,000 elective surgical admissions, ^a age less than 18 years	^a The AHRQ PDI software requires that the hemorrhage or hematoma complicating procedure be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PDI software, the secondary diagnosis could be present on admission. In addition, the control of the hemorrhage or hematoma is not verifiable as following surgery. Consistent with the AHRQ PDI software, the following cases are excluded: obstetric conditions, neonates with a birth weight less than 500 grams, and admissions in which the control of the hemorrhage or hematoma is the only operating room procedure or occurs before the first operating room procedure.
PDI 09	Postoperative respiratory failure per 1,000 elective-surgery admissions, ^a age less than 18 years	^a The AHRQ PDI software requires that the respiratory failure be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PDI software, the secondary diagnosis could be present on admission. In addition, the tracheostomy is not verifiable as following surgery. Consistent with the AHRQ PDI software, the following cases are excluded: admissions with respiratory disease; circulatory disease; craniofacial anomalies with laryngeal or pharyngeal surgery, or with a procedure on face and a diagnosis of craniofacial abnormalities; admissions with a procedure for esophageal resection, lung cancer, or nose, mouth, and pharynx; admissions with degenerative neurological disorders; neuromuscular disorders; neonates with a birth weight less than 500 grams; and admissions in which the tracheostomy is the only operating room procedure.
PDI 10	Postoperative sepsis per 1,000 surgery admissions of length 4 or more days, ^a age less than 18 years	^a The AHRQ PDI software requires that the sepsis be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PDI software, the secondary diagnosis could be present on admission. In addition, the sepsis is not verifiable as following surgery. Consistent with the AHRQ PDI software, the following cases are excluded: admissions with a principal diagnosis of infection; admissions with a procedure for appendicitis, infectious or parasitic diseases, or post-operative infections; obstetric admissions; and neonates.

QI No.	Description	Footnote
PDI 11	Reclosure of postoperative abdominal wound dehiscence per 1,000 abdominopelvic-surgery admissions of length 2 or more days, ^a age less than 18 years	^a Reclosure of abdominal wound dehiscence is not verifiable as following surgery and may have occurred on or before the abdominopelvic procedure. Consistent with the AHRQ PSI software, the following cases are excluded: obstetric admissions; neonates with a birth weight less than 500 grams; admissions in an immunocompromised state or having a procedure code for transplant; admissions with hepatic failure consisting of cirrhosis and hepatic coma or hepatorenal syndrome; and admissions with gasroschisis or umbilical hernia repair in newborns.
PDI 12	Admissions with central venous catheter-related bloodstream infection per 1,000 medical and surgical discharges of length 2 or more days, ^a age less than 18 years	^a The AHRQ PDI software requires that the central venous catheter-related bloodstream infection be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PDI software, the secondary diagnosis could be present on admission. Consistent with the AHRQ PDI software, the following cases are excluded: normal newborns, neonates with a birth weight less than 500 grams, and obstetric admissions.
PDI 14	Admissions for asthma ^a per 100,000 population, ages 2-17	^a Consistent with the AHRQ PDI software, asthma must be the principal diagnosis and the following cases are excluded: admissions with cystic fibrosis or anomalies of the respiratory system, transfers from other institutions, and obstetric admissions.
PDI 15	Admissions with diabetes with short-term complications ^a per 100,000 population, ages 6-17	^a Consistent with the AHRQ PDI software, diabetes must be the principal diagnosis and short-term complications include ketoacidosis, hyperosmolarity, or coma. Transfers from other institutions and obstetric admissions are excluded.
PDI 16	Admissions for pediatric gastroenteritis ^a per 100,000 population, ages 3 months to 17 years	^a Consistent with the AHRQ PDI software, gastroenteritis must be the principal diagnosis or a secondary diagnosis with a principal diagnosis of dehydration. Exclusions include admissions with gastrointestinal abnormalities or bacterial gastroenteritis, transfers from other institutions, neonates if age in days is missing, and obstetric admissions.
PDI 17	Admissions with perforations or abscesses of appendix per 1,000 admissions with appendicitis, ^a ages 1-17	^a Consistent with the AHRQ PDI software, exclusions include transfers from other institutions and obstetric admissions.
PDI 18	Admissions for urinary tract infection (UTI) ^a per 100,000 population, ages 3 months to 17 years	^a Consistent with the AHRQ PDI software, UTI must be the principal diagnosis and the following cases are excluded: kidney or urinary tract disorders, admissions in a immunocompromised state, admissions with hepatic failure consisting of any diagnosis or cirrhosis and hepatic come or hepatorenal syndrome, neonates is age in days is missing, obstetric admissions, and transfers from other institutions.
PDI 90	AHRQ overall Pediatric Quality Indicator (PDI) composite ^a per 100,000 population, ages 6-17	^a Consistent with the AHRQ PDI software, the overall composite is based on the four PDIs for asthma, diabetes, gastroenteritis, and urinary tract infection.
PDI 91	AHRQ acute Pediatric Quality Indicator (PDI) composite ^a per 100,000 population, ages 6-17	^a Consistent with the AHRQ PDI software, the acute composite is based on the two PDIs for gastroenteritis and urinary tract infection.
PDI 92	AHRQ chronic Pediatric Quality Indicator (PDI) composite ^a per 100,000 population, ages 6-17	^a Consistent with the AHRQ PDI software, the chronic composite is based on the two PDIs for asthma and diabetes.

QI No.	Description	Footnote
NQI 01	Admissions with iatrogenic pneumothorax per 1,000 medical and surgical admissions, ^a neonates weighing 500 to 2500 grams	^a The AHRQ PDI software requires that the iatrogenic pneumothorax be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PDI software, the secondary diagnosis could be present on admission. Consistent with the AHRQ PDI software, the following cases are excluded: neonates with a birth weight less than 500 grams, admissions with chest trauma, pleural effusion, thoracic surgery, lung/pleural biopsy, diaphragmatic surgery repair, or cardiac surgery, and obstetric admissions.
NQI 02	Deaths per 1,000 newborn admissions ^a	^a Consistent with the AHRQ PDI software, newborn admissions include babies born outside the hospital and then admitted. Exclusions include newborns weighing less than 500 grams or with any diagnosis of anencephaly, polycystic kidney, trisomy 13, or trisomy 18.
NQI 03	Admissions with blood stream infection per 1,000 medical and surgical discharges of length 2 or more days, newborns ^a	^a The AHRQ PDI software requires that the blood stream infection be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PDI software, the secondary diagnosis could be present on admission. Consistent with the AHRQ PDI software, newborn admissions include babies born outside the hospital and then admitted; infants with a birth weight of 500 to 1499 grams or with gestational age between 24 and 30 weeks; and newborns with a birth weight greater than or equal to 1500 grams, only if the infant experienced death in-hospital, major surgery, mechanical ventilation, or transferred to an acute care facility. Exclusions include newborns weighing less than 500 grams, cases with a principal diagnosis of sepsis or infection or with a length of stay less than 2 days.
Inpatient Quality Indicators¹³		
IQI 8	Deaths per 1,000 hospital admissions with esophageal resection for cancer, ^a age 18 and over	^a Consistent with the AHRQ IQI software, exclusions include obstetric admissions and transfers to another hospital.
IQI 9	Deaths per 1,000 hospital admissions with pancreatic resection for cancer, ^a age 18 and over	^a Consistent with the AHRQ IQI software, exclusions include obstetric admissions and transfers to another hospital.
IQI 11	Deaths per 1,000 hospital admissions with abdominal aortic aneurysm repair, ^a age 18 and over	^a Consistent with the AHRQ IQI software, excluding obstetric admissions and transfers to another hospital.
IQI 12	Deaths per 1,000 hospital admissions with coronary artery bypass graft, ^a age 40 and over	^a Consistent with the AHRQ IQI software, exclusions include obstetric admissions and transfers to another hospital.
IQI 13	Deaths per 1,000 hospital admissions with craniotomy, ^a age 18 and over	^a Consistent with the AHRQ IQI software, exclusions include admissions with a principal diagnosis of head trauma and transfers to another hospital.
IQI 14	Deaths per 1,000 hospital admissions with a hip replacement procedure, ^a age 18 and over	^a Consistent with the AHRQ IQI software, exclusions include hip fractures, obstetric admissions, and transfers to another hospital.
IQI 15	Deaths per 1,000 hospital admissions with acute myocardial infarction (AMI), ^a age 18 and over	^a Consistent with the AHRQ IQI software, AMI must be the principal diagnosis and the following cases are excluded: obstetric admissions and transfers to another hospital.
IQI 16	Deaths per 1,000 hospital admissions with congestive heart failure (CHF), ^a age 18 and over	^a Consistent with the AHRQ IQI software, CHF must be the principal diagnosis and the following cases are excluded: obstetric admissions and transfers to another hospital.

¹³ Indicator IQI 10 is not assigned by the IQI software, version 4. Volume measures IQI 1 to 7 are not calculated.

QI No.	Description	Footnote
IQI 17	Deaths per 1,000 hospital admissions with acute stroke, ^a age 18 and over	^a Consistent with the AHRQ IQI software, stroke must be the principal diagnosis and the following cases are excluded: obstetric admissions and transfers to another hospital.
IQI 18	Deaths per 1,000 hospital admissions with gastrointestinal hemorrhage, ^a age 18 and over	^a Consistent with the AHRQ IQI software, gastrointestinal hemorrhage must be the principal diagnosis and the following cases are excluded: obstetric admissions and transfers to another hospital.
IQI 19	Deaths per 1,000 hospital admissions with hip fracture, ^a age 65 and over	^a Consistent with the AHRQ IQI software, hip fracture must be the principal diagnosis and the following cases are excluded: periprosthetic fractures, obstetric admissions, and transfers to another hospital.
IQI 20	Deaths per 1,000 hospital admissions with pneumonia, ^a age 18 and over	^a Consistent with the AHRQ IQI software, pneumonia must be the principal diagnosis and the following cases are excluded: obstetric admissions and transfers to another hospital.
IQI 21	Cesarean deliveries per 1,000 deliveries ^a	^a Consistent with the AHRQ IQI software, exclusions include deliveries for preterm or multiple infants, deliveries with abnormal presentations or breech procedures, and deliveries resulting in fetal death.
IQI 22	Uncomplicated vaginal birth after cesarean (VBAC) per 1,000 women with previous cesarean deliveries ^a	^a Consistent with the AHRQ IQI software, exclusions include deliveries for preterm or multiple infants, deliveries with abnormal presentations or breech procedures, and deliveries resulting in fetal death.
IQI 23	Laparoscopic cholecystectomies per 1,000 cholecystectomy procedures, ^a age 18 and over	^a Consistent with the AHRQ IQI software, exclusions include obstetric admissions.
IQI 24	Incidental appendectomies per 1,000 hospital admissions with abdominal or pelvic surgery, ^a age 65 and over	^a Consistent with the AHRQ IQI software, exclusions include admissions for cancer involving or adjacent to the appendix, admissions with a colectomy or pelvic evisceration, and obstetric admissions.
IQI 25	Bilateral cardiac catheterizations per 1,000 heart catheterizations for coronary artery disease, ^a age 18 and over	^a Consistent with the AHRQ IQI software, exclusions include valid indications for right-side catheterization and obstetric admissions.
IQI 26	Coronary artery bypass grafts (CABG) ^a per 100,000 population, age 40 and over	^a Consistent with the AHRQ IQI software, exclusions include obstetric admissions.
IQI 27	Percutaneous coronary intervention ^a per 100,000 population, age 40 and over	^a Consistent with the AHRQ IQI software, exclusions include obstetric admissions.
IQI 28	Hysterectomies ^a per 100,000 female population, age 18 and over	^a Consistent with the AHRQ IQI software, exclusions include admissions with genital cancer, pelvic or lower-abdominal trauma, and obstetric admissions.
IQI 29	Laminectomies or spinal fusions ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ IQI software, exclusions include obstetric admissions.
IQI 30	Deaths per 1,000 hospital admissions with percutaneous coronary intervention, ^a age 40 and over	^a Consistent with the AHRQ IQI software, exclusions include obstetric admissions and transfers to another hospital.
IQI 31	Deaths per 1,000 hospital admissions with carotid endarterectomy, ^a age 18 and over	^a Consistent with the AHRQ IQI software, exclusions include obstetric admissions and transfers to another hospital.
IQI 32	Deaths per 1,000 hospital admissions with acute myocardial infarction (AMI), ^a age 18 and over	^a Consistent with the AHRQ IQI software, AMI must be the principal diagnosis and the following cases are excluded: obstetric admissions and transfers to and from another hospital.

QI No.	Description	Footnote
IQI 33	First-time cesarean deliveries (identified by no previous cesarean delivery diagnosis on the record) per 1,000 deliveries ^a	^a Consistent with the AHRQ IQI software, exclusions include previous cesarean delivery, deliveries for preterm or multiple infants, deliveries with abnormal presentations or breech procedures, and deliveries resulting in fetal death.
IQI 34	All vaginal births after cesarean per 1,000 women with previous cesarean deliveries ^a	^a Consistent with the AHRQ IQI software, there are no exclusions.
IQI 90	AHRQ Inpatient Quality Indicator (IQI) mortality composite for selected procedures, ^a age 18 and over	^a Based on seven mortality AHRQ IQIs for esophageal resection for cancer, pancreatic resection for cancer, abdominal aortic aneurysm repair, coronary artery bypass graft (age 40 and over), craniotomy, percutaneous coronary intervention (age 40 and over), and carotid endarterectomy.
IQI 91	AHRQ Inpatient Quality Indicator (IQI) mortality composite for selected conditions, ^a age 18 and over	^a Based on the six mortality AHRQ IQIs for acute myocardial infarction, congestive heart failure, acute stroke, gastrointestinal hemorrhage, hip fracture (age 65 and over), and pneumonia.
Patient Safety Indicators¹⁴		
PSI 2	Deaths per 1,000 hospital admissions with expected low-mortality, ^a age 18 and over or obstetric admissions	^a Consistent with the AHRQ PSI software, admissions with expected low-mortality are identified by Medicare Severity Diagnosis Related Group (MS-DRG) or Diagnosis Related Group (DRG), depending on the date of discharge. Exclusions include admissions with cancer, admissions in an immunocompromised state, admissions involving a traumatic injury, and transfers to an acute care facility.
PSI 4	Deaths per 1,000 elective-surgery admissions having developed specified complications of care during hospitalization, ^a ages 18-89 or obstetric admissions	^a Consistent with the AHRQ PSI software, complications of care include acute renal failure, pneumonia, pulmonary embolism, deep vein thrombosis, sepsis, shock, cardiac arrest, gastrointestinal hemorrhage, and acute ulcer with transfers to another hospital excluded. The AHRQ PSI software requires that the complication be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PSI software, the secondary diagnosis could be present on admission. In addition, the surgery is not verifiable as occurring in the first two days of the inpatient stay.
PSI 6	Admissions with iatrogenic pneumothorax per 1,000 medical and surgical admissions, ^a age 18 and over	^a The AHRQ PSI software requires that the iatrogenic pneumothorax be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PSI software, the secondary diagnosis could be present on admission. Consistent with the AHRQ PSI software, the following cases are excluded: obstetric admissions and admissions with chest trauma, pleural effusion, thoracic surgery, lung/pleural biopsy, diaphragmatic surgery repair, and cardiac surgery.
PSI 7	Admissions with central venous catheter-related bloodstream infection per 1,000 medical and surgical discharges of length 2 or more days, ^a age 18 and over or obstetric admissions	^a The AHRQ PSI software requires that the central venous catheter-related bloodstream infection be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PSI software, the secondary diagnosis could be present on admission. Consistent with the AHRQ PSI software, the following cases are excluded: admissions with a diagnosis of cancer or in an immunocompromised state.

¹⁴ Indicators PSI 1 and 20 are not assigned by the PSI software, version 4. Incidence measures PSI 3 (pressure ulcer), PSI 5 (foreign body), and PSI 16 (transfusion reaction) are not calculated.

QI No.	Description	Footnote
PSI 8	Postoperative hip fracture per 1,000 surgical admissions who were not susceptible to falling, ^a age 18 and over	^a The AHRQ PSI software requires that the hip fracture be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PSI software, the secondary diagnosis could be present on admission. In addition, the hip fracture repair is not verifiable as following surgery. Consistent with the AHRQ PSI software, the following cases are excluded: obstetric cases; admissions for seizure, syncope, stroke, coma, cardiac arrest, poisoning, trauma, delirium and other psychoses, anoxic brain injury, metastatic cancer, lymphoid/bone malignancy malignancy, or self-inflicted injury; admissions for diseases and disorders of the musculoskeletal system and connective tissue; and admissions in which hip fracture repair is the only operating room procedure.
PSI 9	Postoperative hemorrhage or hematoma with surgical drainage or evacuation per 1,000 surgical admissions, ^a age 18 and over	^a The AHRQ PSI software requires that the hemorrhage or hematoma complicating procedure be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PSI software, the secondary diagnosis could be present on admission. In addition, the control of the hemorrhage or hematoma is not verifiable as following surgery. Consistent with the AHRQ PSI software, the following cases are excluded: obstetric conditions and admissions in which the control of the hemorrhage or hematoma is the only operating room procedure.
PSI 10	Postoperative physiologic and metabolic derangements per 1,000 elective-surgery admissions, ^a age 18 and over	^a The AHRQ PSI software requires that the physiologic and metabolic derangements be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PSI software, the secondary diagnosis could be present on admission. In addition, the derangement is not verifiable as following surgery. Consistent with the AHRQ PSI software, the following cases are excluded: obstetric admissions and admissions for ketoacidosis, hyperosmolality, and diabetic coma; admissions with acute renal failure, acute myocardial infarction, cardiac arrhythmia, cardiac arrest, shock, hemorrhage, gastrointestinal hemorrhage, or chronic renal failure.
PSI 11	Postoperative respiratory failure per 1,000 elective-surgery admissions, ^a age 18 and over	^a The AHRQ PSI software requires that the respiratory failure be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PSI software, the secondary diagnosis could be present on admission. In addition, the tracheostomy is not verifiable as following surgery. Consistent with the AHRQ PSI software, the following cases are excluded: admissions with respiratory disease, circulatory disease, craniofacial anomalies, neuromuscular disorders, or degenerative neurological disorder; obstetric admissions; admissions in which the tracheostomy is the only operating room procedure; and admissions with a procedure for esophageal resection, lung cancer, or the nose, mouth, and pharynx.
PSI 12	Postoperative pulmonary embolism (PE) or deep vein thrombosis (DVT) per 1,000 surgical admissions, ^a age 18 and over	^a The AHRQ PSI software requires that the PE or DVT be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PSI software, the secondary diagnosis could be present on admission. The detection of the PE or DVT did not count procedures to unspecified sites. In addition, the interruption of vena cava is not verifiable as following surgery. Consistent with the AHRQ PSI software, the following cases are excluded: obstetric conditions and admissions in which the interruption of vena cava is the only operating room procedure.

QI No.	Description	Footnote
PSI 13	Postoperative sepsis per 1,000 elective-surgery admissions of length 4 or more days, ^a age 18 and over	^a The AHRQ PSI software requires that the sepsis be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PSI software, the secondary diagnosis could be present on admission. In addition, the sepsis is not verifiable as following surgery. Consistent with the AHRQ PSI software, the following cases are excluded: admissions with a principal diagnosis of sepsis or infection, admissions with cancer or in an immunocompromised state; and obstetric admissions.
PSI 14	Reclosure of postoperative abdominal wound dehiscence per 1,000 abdominopelvic-surgery admissions of length 2 or more days, ^a age 18 and over	^a Reclosure of abdominal wound dehiscence is not verifiable as following surgery and may have occurred on or before the abdominopelvic procedure. Consistent with the AHRQ PSI software, the following cases are excluded: obstetric admissions and admissions in an immunocompromised state.
PSI 15	Accidental puncture or laceration during procedure per 1,000 medical and surgical admissions, ^a age 18 and over	^a The AHRQ PSI software requires that the accidental puncture or laceration be reported as a secondary diagnosis (rather than the principal diagnosis), but unlike the AHRQ PSI software, the secondary diagnosis could be present on admission. Consistent with the AHRQ PSI software, the following cases are excluded: obstetric admissions and admissions involving spinal surgery.
PSI 17	Birth trauma - injury to neonate per 1,000 live births ^a	^a Consistent with the AHRQ PSI software, exclusions include newborns weighing less than 2000 grams and newborns with injury to brachial plexus or with osteogenesis imperfecta.
PSI 18	Obstetric trauma ^a per 1,000 instrument-assisted vaginal deliveries	^a Consistent with the AHRQ PSI software, obstetric trauma must involve 3rd or 4th degree lacerations.
PSI 19	Obstetric trauma ^a per 1,000 vaginal deliveries without instrument assistance	^a Consistent with the AHRQ PSI software, obstetric trauma must involve 3rd or 4th degree lacerations.
PSI 21	Admissions for foreign body accidentally left in during procedure ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PSI software, admissions are identified using all diagnosis, including those present on admission.
PSI 22	Admissions for iatrogenic pneumothorax ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PSI software, admissions are identified using all diagnosis, including those present on admission. Exclusions include obstetric admissions and admissions with chest trauma, pleural effusion, thoracic surgery, lung/pleural biopsy, diaphragmatic surgery repair, or cardiac surgery.
PSI 23	Admissions for central venous catheter-related bloodstream infections ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PSI software, admissions are identified using all diagnosis, including those present on admission. Exclusions include admissions with a diagnosis of cancer or in an immunocompromised state.
PSI 24	Admissions for reclosure of abdominal wound dehiscence ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PSI software, admissions are identified using all procedures. Exclusions include obstetric admissions and admissions in an immunocompromised state.
PSI 25	Admissions for accidental puncture or laceration during procedure ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PSI software, admissions are identified using all diagnosis, including those present on admission. Exclusions include obstetric admissions and admissions involving spinal surgery.
PSI 26	Admissions for transfusion reactions ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PSI software, admissions are identified using all procedures.
PSI 27	Admissions for postoperative hemorrhage or hematoma ^a per 100,000 population, age 18 and over	^a Consistent with the AHRQ PSI software, admissions are identified using all procedures. Exclusions include obstetric admissions.

QI No.	Description	Footnote
PSI 90	AHRQ Patient Safety Indicator (PSI) composite, ^a age 18 and over	^a The AHRQ PSI composite has been modified to include the seven PSIs for iatrogenic pneumothorax, central venous catheter-related bloodstream infection, postoperative hip fracture, postoperative pulmonary embolism/deep vein thrombosis, postoperative sepsis, reclosure of postoperative abdominal wound dehiscence, and accidental puncture or laceration. The AHRQ PSI for pressure ulcers is excluded.

Table 2. Sources of 2013 HCUP Inpatient Data for the QDR

Sponsored by the Agency for Healthcare Research and Quality (AHRQ), HCUP is a family of databases, software tools, and products developed through the collaboration of State data organizations, hospital associations, private data organizations, and the Federal government.

HCUP would not be possible without the contributions of the following data collection Partners from across the United States. Use of the HCUP data for the National Healthcare Quality and Disparities Report (QDR) is dependent on when the HCUP Partner joined the project, the availability of the State Inpatient Databases (SID), and the permission from the HCUP Partner to release State-level information in the QDR.

HCUP Partner Organizations
Alaska State Hospital and Nursing Home Association
Arizona Department of Health Services
Arkansas Department of Health
California Office of Statewide Health Planning and Development
Colorado Hospital Association
Connecticut Hospital Association
District of Columbia Hospital Association
Florida Agency for Health Care Administration
Georgia Hospital Association
Hawaii Health Information Corporation
Illinois Department of Public Health
Indiana Hospital Association
Iowa Hospital Association
Kansas Hospital Association
Kentucky Cabinet for Health and Family Services
Louisiana Department of Health and Hospitals
Maine Health Data Organization
Maryland Health Services Cost Review Commission
Massachusetts Center for Health Information and Analysis
Michigan Health & Hospital Association
Minnesota Hospital Association
Mississippi Department of Health
Missouri Hospital Industry Data Institute
Montana MHA – An Association of Montana Health Care Providers
Nebraska Hospital Association
Nevada Department of Health and Human Services

HCUP Partner Organizations
New Hampshire Department of Health & Human Services
New Jersey Department of Health
New Mexico Department of Health
New York State Department of Health
North Carolina Department of Health and Human Services
North Dakota (data provided by the Minnesota Hospital Association)
Ohio Hospital Association
Oklahoma State Department of Health
Oregon Association of Hospitals and Health Systems
Pennsylvania Health Care Cost Containment Council
Rhode Island Department of Health
South Carolina Revenue and Fiscal Affairs Office
South Dakota Association of Healthcare Organizations
Tennessee Hospital Association
Texas Department of State Health Services
Utah Department of Health
Vermont Association of Hospitals and Health Systems
Virginia Health Information
Washington State Department of Health
West Virginia Health Care Authority
Wisconsin Department of Health Services
Wyoming Hospital Association

Table 3. HCUP National and State Inpatient Databases Used for QI Estimates

Use of the HCUP inpatient databases for reporting the AHRQ Quality Indicators (QIs) in the National Healthcare Quality and Disparities Report (QDR) varied by data year and type of reporting.

- **For national QI estimates for data years 2000–2011**, the HCUP Nationwide Inpatient Sample (NIS) was used to calculate national QI estimates for all level of reporting *except by race/ethnicity*. The NIS was not used for reporting QI estimates by race/ethnicity because the availability of race/ethnicity information varied across States and hospitals within States. In addition, the 20 percent sample of the hospitals in the NIS did not provide enough statistical power to detect differences in QI estimates between whites and the other specific racial groups. To facilitate analyses by race/ethnicity, a separate nationally weighted analysis file was constructed from the State Inpatient Databases (SID) and hospitals with good reporting of race/ethnicity using a sampling and weighting strategy similar to the NIS. [Appendix A](#) to this report provides detail on the creation of the analysis files for national estimates.
- **For national QI estimates for data years 2012–2013**, the HCUP National Inpatient Sample (NIS) was not used because the database had been redesigned into a sample of discharges (instead of hospitals) with a revised definition for the target universe that excluded acute long-term care facilities.¹⁵ For consistent estimates before and after data year 2012, nationally weighted analysis files were constructed from the SID using a sampling and weighting strategy similar to the 2000–2011 NIS. In 2012, two analysis file were constructed, one for national estimates not reported by race/ethnicity and a second for reporting by race/ethnicity. In 2013, only one nationally weighted analysis file was created.

The table below lists the HCUP databases used for national QI estimates from 2000–2013, the included SID, and the number of hospitals and unweighted discharges. Use of the SID in the nationally weighted analysis files depended on the availability of the SID when the work began. [Appendix A](#) to this report provides detail on the creation of the analysis files for national estimates.

Year	National inpatient QI estimates, not reported by race/ethnicity	National inpatient QI estimates, reported by race/ethnicity
2000	Nationwide Inpatient Sample (NIS) States (28): AZ, CA, CO, CT, FL, GA, HI, IA, IL, KS, KY, MA, MD, ME, MO, NC, NJ, NY, OR, PA, SC, TN, TX, UT, VA, WA, WI, WV Hospitals: 994 Sample size: 7.5 million records	No analysis file created for this data year

¹⁵ More information on the redesign of the NIS is available on the HCUP user support Web site at https://www.hcup-us.ahrq.gov/db/nation/nis/reports/NIS_2012_Redesign_report.jsp

Year	National inpatient QI estimates, not reported by race/ethnicity	National inpatient QI estimates, reported by race/ethnicity
2001	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (33): AZ, CA, CO, CT, FL, GA, HI, IA, IL, KS, KY, MA, MD, ME, MI, MN, MO, NC, NE, NJ, NY, OR, PA, RI, SC, TN, TX, UT, VA, VT, WA, WI, WV</p> <p>Hospitals: 986</p> <p>Sample size: 7.5 million records</p>	<p>Nationally weighted analysis file with a 20 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (22): AZ, CA, CO, CT, FL, GA, HI, KS, MA, MD, MI, MO, NJ, NY, PA, RI, SC, TN, TX, VA, VT, and WI</p> <p>Hospitals: 976</p> <p>Sample size: 7.7 million discharges</p>
2002	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (35): CA, CO, CT, FL, GA, HI, IA, IL, KS, KY, MA, MD, ME, MI, MN, MO, NC, NE, NJ, NV, NY, OH, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV</p> <p>Hospitals: 995</p> <p>Sample size: 7.9 million records</p>	<p>Nationally weighted analysis file with a 20 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (22): AZ, CA, CO, CT, FL, GA, HI, KS, MA, MD, MI, MO, NJ, NY, PA, RI, SC, TN, TX, VA, VT, and WI</p> <p>Hospitals: 985</p> <p>Sample size: 7.7 million discharges</p>
2003	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (37): AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, MA, MD, MI, MN, MO, NC, NE, NH, NJ, NV, NY, OH, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV</p> <p>Hospitals: 994</p> <p>Sample size: 8.0 million records</p>	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (23): AZ, CA, CO, CT, FL, GA, HI, KS, MA, MD, MI, MO, NH, NJ, NY, PA, RI, SC, TN, TX, VA, VT, and WI</p> <p>Hospitals: 1,711</p> <p>Sample size: 14.6 million discharges</p>
2004	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (37): AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, MA, MD, MI, MN, MO, NC, NE, NH, NJ, NV, NY, OH, OR, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV</p> <p>Hospitals: 1,004</p> <p>Sample size: 8.0 million records</p>	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (23): AR, AZ, CA, CO, CT, FL, GA, HI, KS, MA, MD, MI, MO, NH, NJ, NY, RI, SC, TN, TX, VA, VT, and WI</p> <p>Hospitals: 1,770</p> <p>Sample size: 14.7 million discharges</p>
2005	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (37): AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, MA, MD, MI, MN, MO, NC, NE, NH, NJ, NV, NY, OH, OK, OR, RI, SC, SD, TN, TX, UT, VT, WA, WI, WV</p> <p>Hospitals: 1,054</p> <p>Sample size: 8.0 million records</p>	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (23): AR, AZ, CA, CO, CT, FL, GA, HI, KS, MA, MD, MI, MO, NH, NJ, NY, OK, RI, SC, TN, TX, VT, and WI</p> <p>Hospitals: 1,883</p> <p>Sample size: 15.0 million discharges</p>

Year	National inpatient QI estimates, not reported by race/ethnicity	National inpatient QI estimates, reported by race/ethnicity
2006	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (38): AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, MA, MD, MI, MN, MO, NC, NE, NH, NJ, NV, NY, OH, OK, OR, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV</p> <p>Hospitals: 1,045</p> <p>Sample size: 8.1 million records</p>	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (25): AR, AZ, CA, CO, CT, FL, GA, HI, KS, MA, MD, MI, MO, NH, NJ, NY, OK, RI, SC, TN, TX, UT, VA, VT, and WI</p> <p>Hospitals: 1,908</p> <p>Sample size: 15.0 million discharges</p>
2007	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (40): AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, MA, MD, ME, MI, MN, MO, NC, NE, NH, NJ, NV, NY, OH, OK, OR, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV, WY</p> <p>Hospitals: 1,044</p> <p>Sample size: 8.0 million records</p>	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (26): AR, AZ, CA, CO, CT, FL, GA, HI, KS, MA, MD, MI, MO, NH, NJ, NY, OK, RI, SC, TN, TX, UT, VA, VT, WI, and WY</p> <p>Hospitals: 1,904</p> <p>Sample size: 15.0 million discharges</p>
2008	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (42): AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, NC, NE, NH, NJ, NV, NY, OH, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV, WY</p> <p>Hospitals: 1,056</p> <p>Sample size: 8.2 million records</p>	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (31): AR, AZ, CA, CO, CT, FL, GA, HI, KS, KY, MA, MD, ME, MI, MO, NH, NJ, NV, NY, OK, OR, PA, RI, SC, TN, TX, UT, VA, VT, WI, and WY</p> <p>Hospitals: 1,904</p> <p>Sample size: 15.0 million discharges</p>
2009	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (44): AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MT, NC, NE, NH, NJ, NM, NV, NY, OH, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV, WY</p> <p>Hospitals: 1,050</p> <p>Sample size: 7.8 million records</p>	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (36): AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, KS, KY, MA, MD, ME, MI, MO, NH, NJ, NM, NV, NY, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, and WY</p> <p>Hospitals: 1,967</p> <p>Sample size: 15.7 million discharges</p>

Year	National inpatient QI estimates, not reported by race/ethnicity	National inpatient QI estimates, reported by race/ethnicity
2010	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (45): AK, AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, NE, NJ, NM, NV, NY, OH, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV, WY</p> <p>Hospitals: 1,051</p> <p>Sample size: 7.8 million records</p>	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (37): AK, AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, MA, MD, MI, MO, MS, NC, NJ, NM, NV, NY, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WI, and WY</p> <p>Hospitals: 2,077</p> <p>Sample size: 15.2 million discharges</p>
2011	<p>Nationwide Inpatient Sample (NIS)</p> <p>States (46): AK, AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, ND, NE, NJ, NM, NV, NY, OH, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV, WY</p> <p>Hospitals: 1,049</p> <p>Sample size: 8.0 million records</p>	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (39): AK, AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, MA, MD, ME, MI, MO, MS, NC, NJ, NM, NV, NY, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, and WY</p> <p>Hospitals: 2,076</p> <p>Sample size: 15.4 million discharges</p>
2012	<p>Nationally weighted analysis file with a 20 percent sample of hospitals in the SID</p> <p>States (44): AK, AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, LA, MA, MD, MI, MN, MO, MT, NC, ND, NE, NJ, NM, NV, NY, OH, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV, and WY</p> <p>Hospitals: 1,048</p> <p>Sample size: 7.6 million discharges</p>	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (40): AK, AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, MA, MD, MI, MO, NC, NJ, NM, NV, NY, OH, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, and WY</p> <p>Hospitals: 1,988</p> <p>Sample size: 15.4 million discharges</p>
2013	<p>Nationally weighted analysis file with a 40 percent sample of hospitals in the SID with good reporting of race/ethnicity data</p> <p>States (34): AR, AZ, CA, CO, CT, FL, GA, HI, IA, IL, IN, KS, KY, MD, MI, MO, NC, NJ, NM, NV, NY, OH, OK, OR, PA, SC, SD, TN, TX, VA, VT, WA, WI, and WY</p> <p>Hospitals: 2,063</p> <p>Sample size: 14.9 million discharges</p>	

It should be noted that over time the States include in the NIS and nationally weighted analysis files change as additional statewide data organizations participate in HCUP. Because each file is a sample of hospitals from the States participating in that year (and weighted to the universe of community hospitals nationally), potential exists for different practice patterns across States to influence national measures related to clinical practice over time.

Table 4. Age Groupings for Risk Adjustment

This table shows the 18 categories of patient age, in five-year increments, that are used for risk adjustment. The 36 age-sex categories for risk adjustment are constructed from the 18 age categories split into male and female.

Age Groups
0-4
5-9
10-14
15-17
18-24
25-29
30-34
35-39
40-44
45-49
50-54
55-59
60-64
65-69
70-74
75-79
80-84
85 or older

REFERENCES

- Agency for Healthcare Research and Quality. *Inpatient Quality Indicators: Technical Specifications, AHQR Quality Indicators, Version 4.4 SAS*. Rockville, MD: Agency for Healthcare Research and Quality, March 2012.
- Agency for Healthcare Research and Quality. *Patient Safety Indicators: Technical Specifications, AHQR Quality Indicators, Version 4.4 SAS*. Rockville, MD: Agency for Healthcare Research and Quality, March 2012.
- Agency for Healthcare Research and Quality. *Pediatric Quality Indicators: Technical Specifications, AHQR Quality Indicators, Version 4.4 SAS*. Rockville, MD: Agency for Healthcare Research and Quality, March 2012.
- Agency for Healthcare Research and Quality. *Prevention Quality Indicators: Technical Specifications, AHQR Quality Indicators, Version 4.4 SAS*. Rockville, MD: Agency for Healthcare Research and Quality, March 2012.
- Agency for Healthcare Research and Quality. *Quality Indicator User Guide: Prevention Quality Indicators (PQI) Composite Measures, Version 4.4*. Rockville, MD: Agency for Healthcare Research and Quality, March 2012.
- Coffey R, Barrett M, Houchens R, Moy E, Ho K, Andrews R, Moles E. *Methods Applying AHRQ Quality Indicators to Healthcare Cost and Utilization Project (HCUP) Data for the Ninth (2011) National Healthcare Quality Report (NHQR) and National Healthcare Disparities Report (NHDR)*. HCUP Methods Series Report # 2011-06. Online December 9, 2011. U.S. Agency for Healthcare Research and Quality. Available: <http://www.hcup-us.ahrq.gov/reports/methods.jsp>.
- Coffey R, Barrett M, Houchens R, Moy E, Ho K, Andrews R, Moles E. *Methods Applying AHRQ Quality Indicators to Healthcare Cost and Utilization Project (HCUP) Data for the Seventh (2009) National Healthcare Disparities Report*. HCUP Methods Series Report # 2009-01. Online August 17, 2009. U.S. Agency for Healthcare Research and Quality. Available: <http://www.hcup-us.ahrq.gov/reports/methods.jsp>.
- Fleiss JL. *Statistical Methods for Rates and Proportions*. New York: Wiley, 1973.
- HCUP State Inpatient Databases (SID). Healthcare Cost and Utilization Project (HCUP). 2004, 2001-2013. Agency for Healthcare Research and Quality, Rockville, MD. www.hcup-us.ahrq.gov/sidoverview.jsp.
- HCUP Nationwide Emergency Department Sample (NEDS). Healthcare Cost and Utilization Project (HCUP). 2013. Agency for Healthcare Research and Quality, Rockville, MD. www.hcup-us.ahrq.gov/nedsoverview.jsp.
- HCUP Nationwide Inpatient Sample (NIS). Healthcare Cost and Utilization Project (HCUP). 2000-2011. Agency for Healthcare Research and Quality, Rockville, MD. www.hcup-us.ahrq.gov/nisoverview.jsp.
- Houchens R, Elixhauser A. *Calculating Nationwide Inpatient Sample (NIS) Variances*. HCUP Methods Series Report #2003-2. Revised June 2005. U.S. Agency for Healthcare Research and Quality. Available: <http://www.hcup-us.ahrq.gov/reports/methods.jsp>.

IOM (Institute of Medicine). *Future Directions for the National Healthcare Quality and Disparities Reports*. Washington, DC: The National Academies Press, 2010.

Raetzman S, Stranges E, Coffey RM, Barrett ML, Andrews R, Moy E, Brady J. *Patient Safety in Hospitals in 2004: Toward Understanding Variation Across States*. HCUP Methods Series Report # 2008-02. Online March 14, 2008. U.S. Agency for Healthcare Research and Quality. Available: <http://www.hcup-us.ahrq.gov/reports/methods.jsp>.

The Nielsen Company. *Nielsen Demographic Data*. Available: <http://www.tetrad.com/demographics/usa/nielsen/>

APPENDICES

APPENDIX A: DEVELOPMENT OF NATIONALLY WEIGHTED ANALYSIS FILES FOR THE NATIONAL HEALTHCARE QUALITY AND DISPARITIES REPORT

Nationally weighted analysis files were developed for reporting the AHRQ Quality Indicators (QIs) in the National Healthcare Quality and Disparities Report (QDR) for two reasons:

- The availability of information on patient race/ethnicity varied across States and hospitals within States. In addition, the 20 percent sample of the hospitals in the HCUP Nationwide Inpatient Sample (NIS) did not provide enough statistical power to detect differences in QI estimates between Whites and the other specific racial groups.
- The NIS changed its sampling and weighting strategy in 2012. For consistent national estimates from 2000–2013, weighted analysis files were created for 2012–2013 using the same sampling and weighting strategy employed for the 2000–2011 NIS.

The nationally weighted analysis files were developed using the HCUP State Inpatient Databases (SID). Use of the SID varies by data year and is detailed in [Table 3](#). The nationally weighted analysis files use the same sampling strategy as the 2000–2011 NIS. Hospitals are sampled within five hospital characteristics: geographic region, hospital control (i.e., public, private not-for-profit, and proprietary), urbanized location, teaching status, and size of the hospital based on the number of acute care beds. The 2003–2013 analysis files used for reporting by race/ethnicity used a 40 percent sample of hospitals, instead of a 20 percent sample used in the NIS. The target universe for the nationally weighted analysis files was community hospitals in the United States excluding rehabilitation hospitals. Discharge counts and hospital characteristics for the target universe were obtained from the American Hospital Association Annual Survey Database™. [Table 3](#) lists the number of States, hospitals, and unweighted discharges in all of the nationally weighted analysis files used for the QDR.

For data year 2013, there was only one QDR nationally weighted analysis file developed from 34 SID used for reporting national QI estimates overall and by subpopulations including community income, expected primary payer, and race/ethnicity. In prior years, a separate weighted analysis file was developed for reporting national QI estimates by race/ethnicity.

2013 Nationally Weighted Analysis File for the QDR

Of the 41 SID that were available when the work on the 2015 QDR began, four States did not provide information on patient race to HCUP. One State did not report Hispanic ethnicity. Two States report race and ethnicity, but are missing information on more than half the discharges. The remaining 34 States were used for the creation of the 2013 nationally weighted analysis file. Table A-1 demonstrates the representation by U.S. Census region of these 34 States.

Table A-1. Geographic Representation of the QDR Analysis File, 2013

Census Region	Number of States used for the Nationally Weighted Analysis File	Number of States in the region	Percent of States in the region included in the Analysis File
Northeast	5	9	56%
Midwest	9	12	75%
South	11	16	69%
West	9	13	69%
Total	34	50	68%

Table A-2 compares aggregated totals of various measures for the 34 States as a percent of the national measure. In 2013, the 34 States accounted for 88 percent of U.S. discharges from community, nonrehabilitation hospitals (based on the AHA’s Annual Survey). They accounted for about 85 percent of White and African Americans in the nation and 95 percent of Asian/Pacific Islanders and Hispanics (based on 2013 Nielsen data).

Table A-2. Population Representation of the QDR Analysis File, 2013

Measure	Total for 34 HCUP States with race/ethnicity as a percent of national total
Hospital discharges	88%
Total resident population	87%
<i>Population by race/ethnicity:</i>	
White	85%*
African American	86%*
Asian/Pacific Islander	93%*
Hispanic	95%*
<i>Population by age:</i>	
Population under age 18	87%*
Population age 18-64	87%*
Population over age 64	87%*
<i>Population by income:</i>	
Population with income under the poverty level	85%**

*Calculated using 2013 Nielsen Demographic Update data and 1977 OMB Directive 15 race definitions (e.g. no option for selecting “two or more races”).

**Calculated using Kaiser Family Foundation, statehealthfacts.org. Data Source: Urban Institute and Kaiser Commission on Medicaid and the Uninsured estimates based on the Census Bureau’s March 2014 Current Population Survey (CPS: Annual Social and Economic Supplements), accessed October 23, 2015.

Preparing the 2013 Nationally Weighted QDR Analysis File

The creation of the 2013 nationally weighted analysis file from the SID included the following steps.

1. The sampling frame was defined as community hospitals excluding those that are also rehabilitation hospitals.
2. Hospitals were excluded from the sampling frame if the coding of patient race was suspect (i.e., more than 30 percent of the discharges in the hospital had the race reported as “other”; more than 50 percent of the discharges had no information on the race of the patient; all of the discharges in the hospital had race coded as white, other, or missing; or 100 percent of the discharges had race coded as white and the hospital had more than 50 beds). This caused 4 percent of the hospitals and 2.3 percent of the discharges in 2013 to be excluded. Hospitals were most often excluded because substantial shares of discharges were coded as “other” or “missing” race.
3. A 40-percent stratified sample of hospitals was randomly selected using the strata consistent with the 2000–2011 NIS.
4. For discharges missing race, a “hot deck” imputation method (which draws donors from strata of similar patients within the same hospital) is used to assign values while preserving the variance within the data. Race was missing on 1.8 percent of discharges in the 2013 nationally weighted analysis file.

- Discharge-level weights were developed using the target universe to produce national-level estimates when applied to the analysis file.

The final nationally weighted analysis file in 2013 included about 14.9 million hospital discharges from over 2,000 hospitals.

Evaluating the 2013 Nationally Weighted QDR Analysis File

After creating the 2013 nationally weighted analysis file using the above steps, we evaluated the reliability of national estimates produced with these data by comparing its composition to the 2013 National Inpatient Sample (which was redesigned with a new sample strategy that selects a sample of discharge records from all hospitals participating in HCUP and excludes acute long-term care hospitals). Tables A.3–A.11 contain the distribution of discharges in both files by key demographic and clinical data elements. Based on these analyses, the 2013 nationally weighted analysis file appears to provide reliable national estimates when compared with the 2013 NIS.

Table A-3. Comparison of Census Region, QDR Analysis File and HCUP National Inpatient Sample, 2013

Census Region				
REGION	2013 QDR Analysis File		2013 NIS	
	Frequency	Percent	Frequency	Percent
1: Northeast	7,109,022	19.1	6,730,965	18.9
2: Midwest	8,445,912	22.6	8,004,912	22.5
3: South	14,486,043	38.8	13,818,031	38.8
4: West	7,255,139	19.5	7,043,884	19.8

Table A-4. Comparison of Patient Age, QDR Analysis File and HCUP National Inpatient Sample, 2013

Patient Age in Years at Admission				
AGE	2013 QDR Analysis File		2013 NIS	
	Frequency	Percent	Frequency	Percent
.: Missing	2,074	0.0	1,280	0.0
.A: Invalid	18	0.0	10	0.0
.C: Inconsistent	6,341	0.0	6,635	0.0
0-17	5,659,495	15.2	5,620,416	15.8
18-44	9,094,999	24.4	8,727,809	24.5
45-64	9,277,199	24.9	8,753,270	24.6
65+	13,255,989	35.5	12,488,372	35.1

Table A-5. Comparison of Patient Sex, QDR Analysis File and HCUP National Inpatient Sample, 2013

Patient Sex				
FEMALE	2013 QDR Analysis File		2013 NIS	
	Frequency	Percent	Frequency	Percent
.: Missing	2,737	0.0	4,490	0.0
.A: Invalid	54	0.0	30	0.0
.C: Inconsistent	2,035	0.0	2,720	0.0
0: Male	15,874,986	42.6	15,154,195	42.6
1: Female	21,416,305	57.4	20,436,357	57.4

Table A-6. Comparison of Expected Primary Payer, QDR Analysis File and HCUP National Inpatient Sample, 2013

Expected Primary Payer				
PAY1	2013 QDR Analysis File		2013 NIS	
	Frequency	Percent	Frequency	Percent
.: Missing	45,329	0.1	50,445	0.1
.A: Invalid	7,866	0.0	4,685	0.0
1: Medicare	14,759,370	39.6	13,986,550	39.3
2: Medicaid	7,725,869	20.7	7,417,129	20.8
3: Private Insurance	11,285,820	30.3	10,851,650	30.5
4: Self-pay	2,017,908	5.4	1,862,518	5.2
5: No Charge	210,670	0.6	208,330	0.6
6: Other	1,243,284	3.3	1,216,485	3.4

Table A-7. Comparison of Community Income Quartile, QDR Analysis File and HCUP National Inpatient Sample, 2013

Community Income Quartile Based on the Patient's ZIP Code				
PAY1	2013 QDR Analysis File		2013 NIS	
	Frequency	Percent	Frequency	Percent
.: Missing	824,203	2.2	773,384	2.2
.A: Invalid	58,140	0.2	25,310	0.1
1:First Quartile (lowest income)	11,084,591	29.7	10,199,933	28.7
2:Second Quartile	9,489,423	25.4	9,174,852	25.8
3:Third Quartile	8,432,093	22.6	8,400,391	23.6
4:Fourth Quartile (highest income)	7,407,667	19.9	7,023,922	19.7

Table A-8. Comparison of Patient Race, QDR Analysis File and HCUP National Inpatient Sample, 2013

Patient Race/Ethnicity ¹⁶				
RACE	2013 QDR Analysis File		2013 NIS	
	Frequency	Percent	Frequency	Percent
.: Missing	657,041	1.8	2,235,708	6.3
.A: Invalid	903	0.0	775	0.0
1: White	23,836,610	63.9	22,044,881	61.9
2: Black	5,668,399	15.2	4,951,772	13.9
3: Hispanic	4,740,627	12.7	4,079,463	11.5
4: Asian/Pacific Islander	1,027,776	2.8	940,199	2.6
5: Native American	257,142	0.7	207,740	0.6
6: Other	1,107,618	3.0	1,137,255	3.2

Table A-9. Comparison of Patient Location, QDR Analysis File and HCUP National Inpatient Sample, 2013

Location of Patient Residence				
PL_NCHS	2013 QDR Analysis File		2013 NIS	
	Frequency	Percent	Frequency	Percent
.: Missing	206,031	0.6	173,740	0.5
1: Large central metropolitan	11,134,767	29.9	10,410,481	29.2
2: Large fringe metropolitan	8,958,898	24.0	8,463,155	23.8
3: Medium metropolitan	7,423,607	19.9	6,869,393	19.3
4: Small metropolitan	3,184,217	8.5	3,335,474	9.4
5: Micropolitan (nonmetropolitan)	4,030,039	10.8	3,822,337	10.7
6: Noncore (nonmetropolitan)	2,358,556	6.3	2,523,213	7.1

¹⁶ Differences in race distribution are attributable to higher rates of missing race on the NIS (6%). The 2013 QDR Analysis File uses a modified race variable with missing or invalid values imputed and Native American and Other combined into one racial group.

Table A-10. Comparison of Diagnosis Related Groups (DRG), QDR Analysis File and HCUP National Inpatient Sample, 2013

Top 25 DRGs (Based on QDR Analysis File Frequency)				
MS-DRG, Version 30	2013 QDR Analysis File		2013 NIS	
	Frequency	Percent	Frequency	Percent
795: Normal newborn	2,728,661	7.3	2,624,684	7.4
775: Vaginal delivery w/o complicating diagnoses	2,148,552	5.8	2,083,714	5.9
885: Psychoses	1,144,756	3.1	1,114,212	3.1
470: Major joint replacement or reattachment of lower extremity w/o mcc	1,081,526	2.9	1,025,570	2.9
766: Cesarean section w/o cc/mcc	798,413	2.1	767,080	2.2
871: Septicemia w/o mv 96+ hours w mcc	797,417	2.1	746,425	2.1
392: Esophagitis, gastroent & misc digest disorders w/o mcc	756,207	2.0	718,684	2.0
794: Neonate w other significant problems	690,007	1.9	677,790	1.9
603: Cellulitis w/o mcc	503,628	1.4	478,449	1.3
765: Cesarean section w cc/mcc	487,356	1.3	471,405	1.3
194: Simple pneumonia & pleurisy w cc	424,654	1.1	403,259	1.1
690: Kidney & urinary tract infections w/o mcc	413,751	1.1	390,520	1.1
292: Heart failure & shock w cc	393,008	1.1	369,285	1.0
774: Vaginal delivery w complicating diagnoses	352,468	0.9	342,690	1.0
291: Heart failure & shock w mcc	339,039	0.9	318,960	0.9
641: Nutritional & misc metabolic disorders w/o mcc	336,373	0.9	322,545	0.9
872: Septicemia or severe sepsis w/o mv 96+ hours w/o mcc	335,349	0.9	312,135	0.9
945: Rehabilitation w cc/mcc	322,163	0.9	308,230	0.9
683: Renal failure w cc	320,525	0.9	301,695	0.8
313: Chest pain	311,550	0.8	285,700	0.8
378: GI hemorrhage with cc	295,202	0.8	279,615	0.8
190: Chronic obstructive pulmonary disease w mcc	291,825	0.8	274,100	0.8
897: Alcohol/drug abuse or dependence w/o rehabilitation therapy w/o mcc	287,131	0.8	297,635	0.8
247: Perc cardiovasc proc w drug-eluting stent w/o mcc	284,818	0.8	277,505	0.8
191: Chronic obstructive pulmonary disease w cc	270,011	0.7	253,380	0.7

Abbreviations: cc, complications or comorbidities; mcc, major complications or comorbidities

Table A-11. Comparison of Means, QDR Analysis File and HCUP National Inpatient Sample, 2013

Variable / Label	2013 QDR Analysis File			2013 NIS		
	Minimum	Maximum	Mean	Minimum	Maximum	Mean
LOS: Length of stay (cleaned)	0	365	4.6	0	365	4.5
NDX: Number of diagnoses on this record	0	60	9.5	0	58	9.4
NPR: Number of procedures on this record	0	50	1.7	0	50	1.6
TOTCHG: Total charges (cleaned)	\$100	\$4,994,038	\$40,802	\$100	\$4,991,688	\$39,513

APPENDIX B: MODIFICATION TO THE STATE INPATIENT DATABASES FOR STATE-LEVEL REPORTING BY RACE/ETHNICITY

Data from the each year's State Inpatient Databases (SID) were used to create individual State analysis files that were designed to be used for reporting the State-level Quality Indicators (QIs) in the National Healthcare Quality and Disparities Report (QDR). In SID that include the race/ethnicity of the patient additional data preparation was needed for reporting QI estimates by race/ethnicity. These additional changes to the SID were performed after the modifications described in the HCUP Databases section of this report (e.g., limit the SID to community, nonrehabilitation hospitals, impute missing patient characteristics such as age and sex).

For data year 2013, 35 SID with race/ethnicity were available at the time the work began on the HCUP estimates for the 2015 QDR. Not all of these States participate in reporting State-level QI estimates by race/ethnicity in the QDR. The following steps were taken to further prepare the State-level analysis files for reporting by race/ethnicity:

1. *Exclusion of Hospitals.* We first identified hospitals whose original coding of patient race-ethnicity was "suspect" (i.e., more than 30 percent of the discharges in the hospital had the race reported as "other"; more than 50 percent of the discharges had no information on the race of the patient; all of the discharges in the hospital had race coded as white, other, or missing; or 100 percent of the discharges had race coded as white and the hospital had more than 50 beds).

Table B-1 indicates for data year 2013 the number of excluded hospitals from the race/ethnicity analyses and their associated discharge counts. Twenty-three of the 35 SID had at least one hospital excluded due to suspect race coding. Twelve States had no hospitals with suspect race coding. Overall, 4.0 percent of hospitals and 2.3 percent of discharges were excluded. Except in a few cases, hospitals were most often excluded because substantial shares of discharges were coded as "other" or "missing" race.

Table B-1. Exclusions from State-level Analysis Files for Race/Ethnicity, 2013

Measure	Excluded for any reason	Percent of Total	>30% discharges are "other" race	>50% discharges are "missing" race	All discharges are white, other or missing	All discharges are white and hospital has >50 beds
Total number of hospitals excluded	160	4.0%	74	54	30	2
Total number of discharges excluded	727,757	2.3%	398,565	287,463	41,302	427

2. *Impute for Missing Race/Ethnicity.* Because the area-level QI measures use total State population in the denominator, minimizing the loss of discharges from the numerator for the QI calculation is critical to producing unbiased QI rates. For missing race, we used a "hot deck" imputation method (which draws donors from strata of similar patients within the same hospital) to assign values while preserving the variance within the data. Typically, most hospitals with good reporting of race/ethnicity have no more than five percent of discharges with missing race values before imputation.

3. *Weighting of Selected Hospitals.* We calculated discharge-level weights to account for hospitals excluded because of suspect race coding, community hospitals not reported in the SID, and missing quarters of data. We weighted to the State's universe of hospitals in the American Hospital Association (AHA) Annual Survey Database based on hospital characteristics (teaching status, ownership, urban-rural location, and size of the hospital based on the number of acute care beds).

Some caution should be used in interpreting State comparisons. There may be differences in race and ethnicity coding among States that affect the estimates. For example, some States include Hispanic ethnicity as one of the racial categories, and others record Hispanic ethnicity separately from race. At the hospital-level, policies vary on methods for collecting such data. Some hospitals ask the patient to identify their race and ethnicity, and others determine it from observation. The effect of these and other unmeasured differences in coding of race and ethnicity across the States and hospitals cannot be assessed.

APPENDIX C: EMERGENCY DEPARTMENT RATES FOR SELECTED AHRQ QUALITY INDICATORS

The HCUP Nationwide Emergency Department Samples (NEDS) were used to examine national and regional differences in emergency department (ED) rates for selected AHRQ Prevention Quality Indicators (PQIs) and related Pediatric Quality Indicators (PDIs). The PQIs are measures of quality associated with processes and outcomes of care that occurred in an outpatient or an inpatient setting. The PQIs rely solely on hospital administrative data and, for this reason, are screens for examining quality that may indicate the need for more in-depth studies. Experts have suggested that using both inpatient and emergency room data may give a more accurate picture of avoidable visits/admissions for some ambulatory care sensitive conditions which can be identified by certain PQIs and PDIs. Table C-1 lists the PQIs and PDIs examined.

Table C-1. PQIs and PDIs Used to Examine QI Rates in Inpatient and ED Settings

PQI or PDI	Description
PQI 1	Diabetes with short-term complications
PQI 3	Diabetes with long-term complications
PQI 5	Chronic obstructive pulmonary disease or asthma
PQI 7	Hypertension
PQI 8	Heart failure
PQI 10	Dehydration
PQI 11	Bacterial pneumonia
PQI 12	Urinary tract infections
PQI 13	Angina without cardiac procedure
PQI 14	Uncontrolled diabetes without complications
PQI 15	Adult asthma admissions
PQI 16	Lower extremity amputations among patients with diabetes
PQI 18*	Immunization-preventable influenza
PQI 90	Overall Prevention Quality Indicator (PQI) composite
PQI 91	Acute Prevention Quality Indicator (PQI) composite
PQI 92	Chronic Prevention Quality Indicator (PQI) composite
PDI 14	Pediatric asthma admissions
PDI 15	Pediatric diabetes with short-term complications

* Modified or added version of PQI.

The NEDS contains approximately 30 million ED events from about 950 hospital-based EDs. The NEDS includes information on ED visits that do not result in an admission (i.e., treat-and-release visits and transfers to another hospital) as well as discharge information on patients initially seen in the ED and then admitted to the same hospital. The NEDS was drawn for the State Inpatient Database (SID) and the State Emergency Department Databases (SEDD). Discharge-level weights included with the NEDS are used to produce national estimates.

Table C-2 lists the number of States, EDs, and unweighted ED visits in the NEDS starting in 2008.

Table C-2. States, EDs, and unweighted ED visits in the NEDS

Year	Nationwide Emergency Department Sample (NEDS)
2008	States (28): AZ, CA, CT, FL, GA, HI, IA, IN, KS, KY, MA, MD, ME, MN, MO, NC, NE, NH, NJ, NY, OH, RI, SC, SD, TN, UT, VT, and WI Hospital-based EDs: 980 Sample size: 28.4 million ED visits
2009	States (29): AZ, CA, CT, FL, GA, HI, IA, IN, KS, KY, IL, MA, MD, ME, MN, MO, NC, NE, NH, NJ, NY, OH, RI, SC, SD, TN, UT, VT, and WI Hospital-based EDs: 964 Sample size: 28.9 million ED visits
2010	States (28): AZ, CA, CT, FL, GA, HI, IA, IN, KS, KY, IL, MA, MD, MN, MO, NC, NE, NJ, NV, NY, OH, RI, SC, SD, TN, UT, VT, and WI Hospital-based EDs: 961 Sample size: 28.6 million ED visits
2011	States (30): AZ, CA, CT, FL, GA, HI, IA, IN, KS, KY, IL, MA, MD, ME, MN, MO, NC, ND, NE, NJ, NV, NY, OH, RI, SC, SD, TN, UT, VT, and WI Hospital-based EDs: 951 Sample size: 28.8 million ED visits
2012	States (30): AZ, CA, CT, FL, GA, HI, IA, IN, KS, KY, IL, MA, MD, ME, MN, MO, NC, ND, NE, NJ, NV, NY, OH, RI, SC, SD, TN, UT, VT, and WI Hospital-based EDs: 950 Sample size: 31.1 million ED visits
2013	States (30): AR, AZ, CA, CT, FL, GA, HI, IA, IN, KS, KY, IL, MA, MD, MN, MO, NC, ND, NE, NJ, NV, NY, OH, RI, SC, SD, TN, UT, VT, and WI Hospital-based EDs: 947 Sample size: 28.6 million ED visits

Several steps were taken to prepare the NEDS: (1) QI software review and modification, (2) acquisition of population-based data, (3) handling of missing data, and (4) identification of statistical methods.

1. **QI Software Review and Modification.** A modification of PQI Version 4.4 was used. The PQIs were developed for use with hospital inpatient discharge data. No guidelines for applying the AHRQ QIs to emergency department data were available when this analysis began. Some of the events in the NEDS are visits for patients initially seen in the emergency room and then admitted to the same hospital (an “ED admission”), and some NEDS events are ED visits that do not result in an inpatient admission (e.g., treat-and-release visits and transfers to another hospital). About 15 percent of records in the NEDS represent an ED admission. The PQIs rely on the first-listed diagnosis code (DX1) to identify cases with the outcome of interest. For ED admissions, DX1 is the principal diagnosis code and reflects the condition established to be chiefly responsible for a patients’ admission to the hospital. Unfortunately, principal diagnosis is not clearly discernible for ED visits that do not result in admission. Coding instructions for outpatient data specify that the first-listed diagnosis is supposed to be the “reason for visit,” which is different than the principal diagnosis. Even though DX1 in ED data is not necessarily the principal diagnosis, using DX1 preserves the concept from the PQI algorithm that the first code has higher priority than others. Therefore, this analysis used the first-listed diagnosis in the NEDS.

2. **Acquisition of Population-Based Data.** The next step was to acquire data for the numerator and denominator populations for the PQIs and PDIs in this analysis. These QIs are measures of events that occurs in a hospital, requiring a numerator count of the event of interest and a denominator count of the population (within the geographic area) to which the event relates.

For the numerator counts of the PQI or PDI, we used the HCUP NEDS to create national estimates of all ED visits, ED visits resulting in admission to the same hospital, and all other types of ED visits. For the denominator counts, population ZIP-Code-level counts from demographic update data provided by Nielsen (a vendor that compiles and adds value to the U.S. Bureau of Census data) were used for all reporting categories. Nielsen uses intra-census methods to estimate household and demographic statistics for geographic areas (The Nielsen Company, 2012). We also used the Nielsen population data for risk adjustment by age and sex.

3. **Preparation of HCUP Data.** Next, “hot deck” imputation method (which draws donors from strata of similar hospitals and patients) was used to assign missing values of patient age and sex. Patient age and sex are missing on less than 0.02 percent of the records in the NEDS.

4. **Statistical Methods.** Age-sex adjustments were made for age and sex differences across population subgroups and were based on methods of direct standardization (Fleiss, 1973). Age was categorized into 18 five-year increments. Statistical methods are explained in more detail in [Appendix D](#) to this report.

5. **Masking Rates for Statistical Reliability, Data Quality, and Confidentiality.** PQI and PDI estimates were included in this analysis if they reached a threshold defined by a relative standard error less than 30 percent and at least 11 unweighted cases in the

denominator. Estimates that did not meet this threshold were suppressed and the corresponding table cell was marked with an asterisk.

APPENDIX D: STATISTICAL METHODS

This appendix explains the statistical methods and gives formulas for the calculations of standard errors and hypothesis tests. These statistics are derived from multiple databases: the NIS, the NEDS, the nationally weighted analysis files, the SID, and demographic population data provided by Nielsen (a vendor that compiles and adds value to Bureau of Census data). For NIS, NEDS, and nationally weighted analysis files, the standard errors are calculated as described in the HCUP report entitled *Calculating Nationwide Inpatient Sample (NIS) Variances* (Houchens, et al., 2005). We will refer to this report simply as the NIS Variance Report throughout this appendix. This method takes into account the cluster and stratification aspects of the sample design when calculating these statistics using the SAS procedure PROC SURVEYMEANS. For the SID we used the same procedure omitting the cluster and stratification features. For population counts based on Nielsen data, there is no sampling error.

Even though the SID databases contain nearly all discharges from nearly all hospitals in the State, we treat the files as though they were drawn from an infinite population. We do not employ finite population correction factors in estimating standard errors. We take this approach because we view the outcomes as a result of myriad processes that go into treatment decisions rather than being the result of specific, fixed processes generating outcomes for a specific population and a specific year. We consider the SID to be samples from a “super-population” for purposes of variance estimation. Further, we assume the counts (of QI events) to be binomial.

1. Area-Based QIs using Weighted Discharge Data in the Numerator and Nielsen Population Data in the Denominator

a. Standard error estimates for discharge rates per 100,000 population using the 2013 Nielsen population data.

The observed rate was calculated as follows:

$$R = 100,000 \cdot \frac{\sum_{i=1}^n w_i x_i}{N} = 100,000 \cdot \frac{S}{N}. \quad (\text{D.1})$$

w_i and x_i , respectively, are the weight and variable of interest for patient i in the quality analysis file or SID. To obtain the estimate of S and its standard error, SE_S , we followed instructions in the NIS Variance Report (modified for the SID, as explained above)

The population count in the denominator is a constant. Consequently, the standard error of the rate R was calculated as:

$$SE_R = 100,000 \cdot SE_S / N. \quad (\text{D.2})$$

b. Standard error estimates for age/sex adjusted inpatient rates per 100,000 population using the Nielsen population data.

We adjusted rates for age and sex using the method of direct standardization (Fleiss, 1973). We estimated the observed rates for each of 36 age/sex categories (described in [Table 4](#) in this methods report, Age Groupings for Risk Adjustment). We then calculated the weighted average of those 36 rates using weights proportional to the percentage of a standard population in each cell. Therefore, the adjusted rate represents the rate that would be expected for the observed study population if it had the same age and sex distribution as the standard population.

For the standard population we used the age and sex distribution of the U.S. as a whole according to the year 2010. In theory, differences among adjusted rates were not attributable to differences in the age and sex distributions among the comparison groups because the rates were all calculated with a common age and sex distribution.

The adjusted rate was calculated as follows (and subsequently multiplied by 100,000):

$$A = \frac{\sum_{g=1}^{36} N_{g,std} \sum_{i=1}^{n(g)} \frac{w_{g,i} x_{g,i}}{N_{g,obs}}}{\sum_{g=1}^{36} N_{g,std}} = \frac{\sum_{g=1}^{36} \sum_{i=1}^{n(g)} \frac{N_{g,std}}{N_{g,obs}} w_{g,i} x_{g,i}}{N_{std}} = \frac{\sum_{g=1}^{36} \sum_{i=1}^{n(g)} w_{g,i}^* x_{g,i}}{N_{std}} = \frac{S^*}{N_{std}} \quad (D.3)$$

g = index for the 36 age/sex cells.

$N_{g,std}$ = Standard population for cell g (year 2010 total US population in cell g).

$N_{g,obs}$ = Observed population for cell g (year 2013 subpopulation in cell g , e.g., females, state of California, etc.).

$n(g)$ = Number in the sample for cell g .

$x_{g,i}$ = Observed quality indicator for observation i in cell g (e.g., 0 or 1 indicator).

$w_{g,i}$ = Quality analysis file or SID discharge weight for observation i in cell g .

The estimates for the numerator, S^* , and its standard error, SE_{S^*} , were calculated in similar fashion to the unadjusted estimates for the numerator S in formula A.1. The only difference was that the weight for patient i in cell g was redefined as:

$$w_{g,i}^* = \frac{N_{g,std}}{N_{g,obs}} \cdot w_{g,i} \quad (D.4)$$

Following instructions in the NIS Variance Report (modified for the SID, as explained above), we used PROC SURVEYMEANS to obtain the estimate of S^* , the weighted sum in the numerator using the revised weights, and the estimate SE_{S^*} , the standard error of the weighted sum S^* . The denominator is a constant. Therefore, the standard error of the adjusted rate, A , was calculated as

$$SE_A = 100,000 SE_{S^*} / N_{std}. \quad (D.5)$$

2. Provider-based QIs using Weighted Discharge Data in the Numerator and Denominator

a. Standard error estimates for inpatient rates per 1,000 discharges using discharge counts in both the numerator and the denominator.

We calculated the observed rate as follows:

$$R = 1,000 \cdot \frac{\sum_{i=1}^n w_i x_i}{\sum_{i=1}^n w_i} = 1,000 \cdot \frac{S}{N}. \quad (D.6)$$

Following instructions in the HCUP NIS Variance Report (modified for the SID, as explained above), we used PROC SURVEYMEANS to obtain estimates of the weighted mean, S/N , and the standard error of the weighted mean, $SE_{S/N}$. We multiplied this standard error by 1,000.

b. Standard error estimates for age/sex adjusted inpatient rates per 1,000 discharges using inpatient counts in both the numerator and the denominator.

We used the 2010 NIS national estimates for the standard inpatient population age-sex distribution. For each of the 36 age-sex categories, we estimated the number of U.S. inpatient discharges, $\hat{N}_{g,std}$, in category g . We calculated the directly adjusted rate:

$$A = 1,000 \cdot \frac{\sum_{g=1}^{36} \hat{N}_{g,std} \frac{\sum_{i=1}^{n(g)} w_{g,i} x_{g,i}}{n(g)}}{\sum_{g=1}^{36} \hat{N}_{g,std}} = 1,000 \cdot \sum_{g=1}^{36} \hat{P}_{g,std} \frac{\sum_{i=1}^{n(g)} w_{g,i} x_{g,i}}{\sum_{i=1}^{n(g)} w_{g,i}}. \quad (D.7)$$

g = index for the 36 age/sex cells.

$\hat{N}_{g,std}$ = Standard inpatient population for cell g (Estimate of year 2010 total inpatient population for cell g).

$n(g)$ = Number in the sample for cell g .

$x_{g,i}$ = Observed quality indicator for observation i in cell g .

$w_{g,i}$ = Quality analysis file or SID discharge weight for observation i in cell g .

Note that $\hat{P}_{g,std} = \frac{\hat{N}_{g,std}}{\sum_{g=1}^{36} \hat{N}_{g,std}}$ is the proportion of the standard inpatient population in cell g .

Consequently, the adjusted rate is a weighted average of the cell-specific rates with cell weights equal to $\hat{P}_{g,std}$. These cell weights are merely a convenient, reasonable standard

inpatient population distribution for the direct standardization. Therefore, we treat these cell weights as constants in the variance calculations:

$$SE(A) = \sqrt{Var(A)} = 1,000 \cdot \sqrt{Var \left(\sum_{g=1}^{36} \hat{P}_{g,std} \frac{\sum_{i=1}^{n(g)} w_{g,i} X_{g,i}}{\sum_{i=1}^{n(g)} w_{g,i}} \right)} = 1,000 \cdot \sqrt{\sum_{g=1}^{36} \hat{P}_{g,std}^2 \cdot Var \left(\frac{\sum_{i=1}^{n(g)} w_{g,i} X_{g,i}}{\sum_{i=1}^{n(g)} w_{g,i}} \right)}. \quad (A.8)$$

The variance of the ratio enclosed in parentheses was estimated separately for each cell g by squaring the SE calculated using the method of section 2.a:

$$SE(A) = 1,000 \cdot \sqrt{\sum_{g=1}^{36} \hat{P}_{g,std}^2 \cdot \{SE(R_g)\}^2}$$

$$R_g = \frac{\sum_{i=1}^{n(g)} w_{g,i} X_{g,i}}{\sum_{i=1}^{n(g)} w_{g,i}} \quad (A.9)$$

Following instructions in the HCUP NIS Variance Report (modified for the SID, as explained above), we used PROC SURVEYMEANS to obtain estimates of the weighted means, R_g , and their standard errors.

3. Significance tests.

Let R_1 and R_2 be either observed or adjusted rates calculated for comparison groups 1 and 2, respectively. Let SE_1 and SE_2 be the corresponding standard errors for the two rates. We calculated the test statistic and (two-sided) p-value:

$$t = \frac{R_1 - R_2}{\sqrt{SE_1^2 + SE_2^2}} \quad (A.10)$$

$$p = 2 * \text{Prob}(Z > |t|)$$

where Z is a standard normal variate.

Note: the following functions calculate p in SAS and EXCEL:

SAS: $p = 2 * (1 - \text{PROBNORM}(\text{ABS}(t)))$;

EXCEL: $= 2*(1 - \text{NORMDIST}(\text{ABS}(t), 0, 1, \text{TRUE}))$

APPENDIX E: CAVEATS TO THE INTERPRETATION OF HCUP-BASED QI ESTIMATES REPORTED IN THE NATIONAL HEALTHCARE QUALITY AND DISPARITIES REPORT

Some caution should be used in interpreting the AHRQ Quality Indicators (QI) statistics presented in the National Healthcare Quality and Disparities Report (QDR). These caveats relate to the how the AHRQ QIs were applied to the HCUP data, ICD-9-CM coding changes, inter-State differences in data collection, and other more general issues.

ICD-9-CM Coding Changes: A number of the AHRQ QIs are based on diagnoses and procedures for which ICD-9-CM coding has generally become more specific over the period of this study. If coding changes cause earlier estimates to be non-comparable to the later estimates, then the earlier estimates are not reported. For this reason, the following QIs are not reported for certain years:

- *Not reported prior to 2004:* birth trauma (PSI 17)
- *Not reported prior to 2005:* the PQI for chronic obstructive pulmonary disease (PQI 5), the overall PQI composite (PQI 90), and chronic PQI composite (PQI 92), the PSI for deaths per elective-surgery admissions having developed specified complications of care during hospitalization (PSI 4), and the PSI for post-operative pulmonary embolism or deep vein thrombosis (PSI 12).
- *Not reported prior to 2008:* QIs for sepsis (PDI 10 and PSI 13) and blood stream infections (NQI 3, PDI 12, PSI 7, and PSI 23).
- *Not reported for any year:* QIs for pressure ulcer (PDI 2 and PSI 3) because numerous coding changes between 2000 and 2012 make longitudinal analysis inadvisable.

Data Collection Differences Among States: Organizations providing statewide data generally collect the data using the Uniform Billing format (UB-04) and, for earlier years, the UB-92 or Uniform Hospital Discharge Data Set (UHDDS) format. However, not every statewide data organization collects all data elements nor codes them the same way. For the QDR, uneven availability of a few data elements underlie some estimates, as noted next.

Data Elements Used for QI Exclusions: Three data elements required for certain QIs were not available in every State: “secondary procedure day,” “admission type” (elective, urgent, newborn, and emergency), and “present on admission.” We modified the AHRQ QI software in instances where these data elements are used to exclude specific cases from the QI measures:

- Some of the PSIs and PDIs use procedure days to determine the timing of a patient safety event. In States without procedure days, the patient safety event cannot be verified as following surgery.¹⁷ Seven PSIs and three PDIs use procedure day to qualify safety events as post-operative, excluding patients for which the event is not pre-operative:
 - PSI 8 – Post-operative hip fractures
 - PSI 9 – Post-operative hemorrhage or hematoma
 - PSI 10 – Post-operative physiologic/metabolic derangements
 - PSI 11 – Post-operative respiratory failure
 - PSI 12 – Post-operative pulmonary embolism or deep vein thrombosis

¹⁷ Several States are missing data on day of procedure. The states without procedure days in the 2004-2013 SID include: Oklahoma, Utah, and West Virginia. For 2004-2011, Ohio did not have procedure days. For 2004, Illinois, Kansas, and Washington did not have procedure days.

- PSI 14 – Post-operative abdominal wound dehiscence
 - PSI 27 – Post-operative hemorrhage or hematoma (area based)
 - PDI 8 – Pediatric: Post-operative hemorrhage or hematoma
 - PDI 9 – Pediatric: Post-operative respiratory failure
 - PDI 11 – Pediatric: Post-operative wound dehiscence
- PSI 4 “Deaths per 1,000 elective-surgery admissions having developed specified complications of care during hospitalization” uses the day of the principal procedure or type of admission to identify an elective admission. All of the States that do not report day of principal procedure do report type of admission, so one of the other can be used.
 - For QIs that use admission type “elective” and “newborn,” we imputed the missing admission type using available information. For all States except California, an admission type of “elective” was assigned if the DRG did not indicate trauma, delivery, or newborn. An admission type of “newborn” was assigned if the DRG indicated a newborn. For California, which did not provide any information on admission type, information on scheduled admissions was used to identify elective admissions and DRGs were used to identify newborn admissions.
 - For QIs that use present on admission (POA), we modified the AHRQ QI software to calculate indicators without considering whether the condition was present at admission. Ten PSIs and 9 PDIs used information on whether a condition was present on admission (POA) to exclude patients:
 - PSI 6 Iatrogenic Pneumothorax
 - PSI 7 Central Venous Catheter-Related Bloodstream Infection
 - PSI 8 Postoperative Hip Fracture
 - PSI 9 Postoperative Hemorrhage or Hematoma
 - PSI 10 Postoperative Physiologic and Metabolic Derangements
 - PSI 11 Postoperative Respiratory Failure
 - PSI 12 Postoperative Pulmonary Embolism or Deep Vein Thrombosis
 - PSI 13 Postoperative Sepsis
 - PSI 14 Postoperative Abdominal Wound Dehiscence (Provider-based)
 - PSI 15 Accidental Puncture or Laceration (Provider-based)
 - PDI 1 Pediatric: Accidental Puncture or Laceration
 - PDI 5 Pediatric: Iatrogenic Pneumothorax
 - PDI 8 Pediatric: Postoperative Hemorrhage or Hematoma
 - PDI 9 Pediatric: Postoperative Respiratory Failure
 - PDI 10 Pediatric: Postoperative Sepsis
 - PDI 11 Pediatric: Postoperative Abdominal Wound Dehiscence
 - PDI 12 Pediatric: Central Venous Catheter-Related Bloodstream Infection
 - NQI 01 Neonatal Iatrogenic Pneumothorax
 - NQI 03 Neonatal Central Venous Catheter-Related Bloodstream Infection

Number of Clinical Fields: Another data collection issue relates to the number of fields that statewide data organizations permit for reporting patients’ diagnoses and procedures during the hospitalization. The SID for different States can contain as few as 6 or as many as 50 or more fields for reporting diagnoses and procedures, as shown in Table E-1. The more fields used, the more quality-related events that can be captured. However, in an analysis of year 2000 data, 95 percent of discharge records captured all of patients’ diagnoses in 10 to 13 data elements in States with 30 diagnosis fields available. For States with 30 procedure fields

available, 95 percent of records captured all of patients' procedures in five fields. Thus, limited numbers of fields available for reporting diagnoses and procedures are unlikely to have much effect on results, because all statewide data organizations participating in HCUP allow at least nine diagnoses and six procedures. All available diagnosis and procedure fields in the SID were used for the QDR analyses, so that the full richness of the databases would be included.

Table E-1. Number of Diagnosis and Procedure Fields by State, 2013

State	Number of Diagnoses	Number of Procedures
Arkansas	19	8
Arizona	26	12
California	25	21
Colorado	30	30
Connecticut	31	30
District of Columbia	25	24
Florida	32	31
Georgia	50	50
Hawaii	25	25
Illinois	26	25
Indiana	61	25
Iowa	51	55
Kansas	30	25
Kentucky	26	25
Louisiana	10	6
Maryland	30	30
Massachusetts	15	15
Michigan	31	30
Minnesota	51	50
Missouri	30	25
Montana	26	25
Nebraska	10	6
Nevada	34	25
New Jersey	25	25
New Mexico	19	6
New York	26	14
North Carolina	26	20
North Dakota	47	25
Ohio	17	9
Oklahoma	17	16
Oregon	26	25
Pennsylvania	19	6
Rhode Island	26	25
South Carolina	16	13
South Dakota	26	25
Tennessee	19	6
Texas	26	15
Utah	9	6
Vermont	21	20
Virginia	18	6
Washington	25	25

State	Number of Diagnoses	Number of Procedures
West Virginia	19	6
Wisconsin	51	50
Wyoming	30	25

Use of External Cause-of-Injury Codes: Another issue relates to the reporting of external cause-of-injury codes (E Codes). Five of the 25 PSIs and one of the PDIs use E codes to help identify complications of care or to exclude cases (e.g., poisonings, self-inflicted injury, and trauma) from numerators and denominators, as shown in Table E-2.

Table E-2. Use of E codes in the AHRQ Quality Indicators, Version 4.4

PSI or PDI *	Codes used for defining the numerator		Codes used for defining exclusions	
	E codes	Similar ICD-9-CM codes	E codes	Similar ICD-9-CM codes
PSI 21	E8710 – E8719	9984, 9987	None	None
PSI 8	None	None	Self-inflicted injury (E95nn); Poisoning (E85nn, E86nn, E951n, E952n, E962nn, E980n-E982n)	9600-9799
PSI 15 PSI 25 PDI 1	E870n	9982	None	None
PSI 26	E8760	9996-9997	None	None

* All other PSIs and PDIs do not use E codes.

The E codes in the AHRQ PSI and PDI software have been augmented wherever possible with similar ICD-9-CM codes. Uneven capture of these data has the potential of affecting rates and should be kept in mind when judging the level of these events.

While all HCUP States report E Codes, the policies on reporting medical misadventures and adverse effects can vary:

- California (through data year 2009) and Washington do not require hospitals to report E codes in the range E870-E879 (medical misadventures and abnormal reactions).
- Georgia does not report E codes in the range E870-E879 (medical misadventures and abnormal reactions) and E930-E949 (adverse effects).
- South Carolina (through data year 2007) did not report E codes in the range E870-E876 (medical misadventures).

Non-Resident Discharges in State-Level Estimates: HCUP databases include discharges from all hospitals in a State, and may include non-residents, including foreign patients, which can bias the results for QIs using area-based denominators (State populations). We had no way to adjust the HCUP data to consistently exclude the non-resident discharges and include discharges for residents hospitalized in other States. Therefore, non-resident discharges were

retained in the SID databases for the QDR analyses. Based on an analysis performed with the 2013 SID, the overall percentage of non-resident discharges within a State is 4 percent with a range from 1 percent to 45 percent. Most States were below 10 percent, but five States (ND, SD, TN, VT, WV) and the District of Columbia had more than 10 percent of discharges in the SID that were for non-residents.

Variation Among State QI Rates: Variation in State rates can be caused by many factors, including differences in practice patterns, underlying disease prevalence, health behaviors, access to health insurance, income levels of the population, demographics, spending on health services, supply of health care resources, coding conventions, and so on. To understand some of the variation in State rates, we analyzed the 2001 State-level QI rates in relation to these types of factors. For more information on this study, refer to the *Methods Applying AHRQ Quality Indicators to the Healthcare Cost and Utilization Project (HCUP) Data for the Ninth (2011) NHQR and NHDR* (Coffey et al., 2011). The report includes an appendix that describes analyses performed for each Prevention Quality Indicator (PQI), and the result in terms of whether the factors (with each tested separately because of the limited number of observations) were positively, negatively, or not significantly related to the QIs.

In a subsequent analysis, we investigated sources of variation in Patient Safety Indicator (PSI) rates across States using 2004 data. The analysis concluded there were few state factors (such as state policy, hospital characteristics, coding practices, and socio-demographics) with strong patterns of association to State-level variation in the nine PSI rates studied. The strongest result occurred with coding practices — the number of diagnosis fields coded. Only one in five correlations between the PSIs and State factors were statistically significant, although there is generally no pattern. For more information on this study, refer to the *Methods Applying AHRQ Quality Indicators to the Healthcare Cost and Utilization Project (HCUP) Data for the Ninth (2011) NHQR and NHDR* (Coffey et al., 2011). The report includes the executive summary from the report, *Patient Safety in Hospitals in 2004: Toward Understanding Variation Across States*.

APPENDIX F: QDR SUMMARY MEASURES FOR PATIENT SAFETY AND MORTALITY FOR SELECTED PROCEDURES AND CONDITIONS

To examine national and State-level trends in inpatient mortality and patient safety events, risk-adjusted rates for select AHRQ Inpatient Quality Indicators (IQIs) and Patient Safety Indicators (PSIs) were summarized. The three QDR summary measures include the following:

1. Mortality for selected conditions based on select IQIs
2. Mortality for selected procedures based on select IQIs
3. Patient safety based on select PSIs

These summary measures were calculated as a weighted sum of risk-adjusted rates for individual IQIs and PSIs. The weights used to calculate the QDR summary measures were relatively consistent with AHRQ IQI and PSI Composites; however, the methodology employed to perform the calculations differed. The IQI and PSI composites were designed for use with hospital-level rates, while the QDR report only national and State-level statistics.

IQIs and PSIs Used for the QDR Summary Measures

The QDR summary measure for mortality for selected conditions was based on six IQIs also included in the similar IQI Composite (Table F-1). The IQI composite weights were extracted from the SAS software, version 4.4. They are based on pooled SID denominators (i.e., the relative frequency of the denominators of the component indicators). This approach is known as “opportunity weighting,” because it gives equal weight to each opportunity that the health care system has to do “the right thing,” which in this case is to discharge the patient alive from the hospital. The QDR summary measure weights were the same as the weights in the similar IQI Composite.

Table F-1. IQIs Included in the QDR Summary Measure for Mortality for Selected Conditions

IQI	Description	IQI Composite Weight	QDR Summary Measure Weight
IQI 15	Acute Myocardial Infarction	0.1433	0.1433
IQI 16	Congestive Heart Failure	0.2739	0.2739
IQI 17	Acute Stroke Adult Mortality Rate	0.1329	0.1329
IQI 18	Gastrointestinal Hemorrhage	0.1302	0.1302
IQI 19	Hip Fracture	0.0678	0.0678
IQI 20	Pneumonia	0.2519	0.2519

The QDR summary measure for mortality for selected procedures was based on four IQIs instead of the eight IQIs included in the similar IQI Composite (Table F-2). Three IQIs were excluded because the procedures were not high-volume at the State level and, therefore, State-level risk-adjusted rates were often unavailable. The IQI for Hip Replacement has a zero-weight in the IQI Composite because it was not endorsed by the National Quality Forum (NQF) (for version 4.4 IQIs). The IQI composite weights were extracted from the SAS software, version 4.4, and were also based on pooled SID denominators. The IQI Composite weights were proportionally reallocated into the QDR summary measure weights to account for the excluded IQIs.

Table F-2. IQIs Included in the QDR Summary Measure for Mortality for Selected Procedures

IQI	Description	IQI Composite Weight	QDR Summary Measure Weight
IQIs Included in the QDR Summary			
IQI30	Percutaneous coronary intervention	0.5659	0.6275
IQI12	CABG	0.2001	0.2219
IQI13	Craniotomy	0.1031	0.1143
IQI11	Abdominal Aortic Aneurysm Repair	0.0328	0.0364
IQIs Excluded in the QDR Summary, but in the IQI Composite			
IQI08	Esophageal Resection	0.0043	0.0000
IQI09	Pancreatic Resection	0.0048	0.0000
IQI14	Hip Replacement	0.0000	0.0000
IQI31	Carotid Endarterectomy	0.0890	0.0000

The QDR summary measure for patient safety was based on seven PSIs instead of the eleven PSIs included in the similar PSI Composite (Table F-3). One PSI Pressure Ulcer was excluded due to its dependence upon reporting whether the diagnosis is present on admission (POA) to the hospital. (This information is not uniformly available across HCUP States). Three PSIs have zero weights in the PSI Composite because they were not endorsed by the NQF (for version 4.4 PSIs). The PSI composite weights were extracted from the SAS software, version 4.4, and are based on pooled SID numerators (i.e., the relative frequency of the numerators of the component indicators). This approach is known as “event weighting,” because it gives equal weight to each event, regardless of how many patients were evaluated for the occurrence of that event. The PSI Composite weights were proportionally reallocated into the QDR summary measure weights to account for the excluded PSIs.

Table F-3. PSIs Included in the QDR Summary Measure for Patient Safety

PSI	Description	PSI Composite Weight	QDR Summary Measure Weight
PSIs Included in the QDR Summary			
PSI15	Accidental Puncture or Laceration	0.2982	0.3925
PSI12	Postoperative Pulmonary Embolism or Deep Vein Thrombosis	0.2360	0.3106
PSI07	Central Venous Catheter-Related Bloodstream Infections (2008 only)	0.1280	0.1685
PSI06	Iatrogenic Pneumothorax	0.0457	0.0602
PSI13	Postoperative Sepsis (2008 only)	0.0383	0.0504
PSI14	Postoperative Wound Dehiscence	0.0124	0.0163
PSI08	Postoperative Hip Fracture	0.0011	0.0014

PSI	Description	PSI Composite Weight	QDR Summary Measure Weight
PSIs Excluded in the QDR Summary, but in the PSI Composite			
PSI03	Pressure Ulcer	0.2403	0.0000
PSI09	Postoperative Hemorrhage or Hematoma	0.0000	0.0000
PSI10	Postoperative Physiologic and Metabolic Derangement	0.0000	0.0000
PSI11	Postoperative Respiratory Failure	0.0000	0.0000

Calculation of the IQI and PSI Summary Measures

Each summary measure was calculated as follows:

$$s = \sum_i a_i X_i$$

Where a_i corresponds to the weight to the i^{th} QI and X_i corresponds to the risk-adjusted rate for the i^{th} QI.

The standard error (SE) of the summary measure is the square-root of the variance:

$$\text{Var} \left(\sum_i a_i X_i \right) = \sum_i a_i^2 \text{Var}(X_i) + \sum_i \sum_{j \neq i} a_i a_j \text{Cov}(X_i, X_j)$$

Where a_i corresponds to the weight to the i^{th} QI and X_i corresponds to the risk-adjusted rate for the i^{th} QI. The correlations actually had very little effect on the estimated SE for the summary measures. For example, in examining mortality for select conditions, the SE was 0.293 if we assume the correlations are zero (i.e., the individual measures are uncorrelated) and the SE was 0.303 if we assume the correlations are those estimated by the covariance matrix of the State-level rates, which were in the range of 70 to 85 percent. Therefore, the SEs were calculated on the assumption that the individual measures were independent of one another, which eliminates the second term on the right-hand side of the formula above.